# 2393-199

10-28-6

## 1970

## DIAZINON® SEED PROTECTANT

TO PROTECT GERMINATING CORN SEEDS AND SEEDLINGS AGAINST SEED CORN BEETLES AND SEED CORN MAGGOT.

THIS PRODUCT CONTAINS GRAPHITE TO ASSIST IN ACCURATE PLANTING.

| ACTIVE INGREDIENT:         |     |      |      |     |      |     |      |     |     |    |       |
|----------------------------|-----|------|------|-----|------|-----|------|-----|-----|----|-------|
| O,O-diethyl O-(2-isopre    | ору | /1-4 | -m   | eth | yl-é | g-S | vri: | nid | inv | 4) |       |
| phosphorothioate           |     |      |      |     |      |     |      |     |     |    | . 33% |
| INERT INGREDIENTS          |     |      |      |     |      |     |      |     |     |    | . 67% |
| * Trademark Geigy Chemical | Col | rno  | rati | on  |      |     |      |     |     |    | 100%  |

### WARNING

#### KEEP OUT OF REACH OF CHILDREN

See side panels for complete warning statement, use instructions, and notice of warranty.

**NET WEIGHT** 

ONE POUND

**MEASURING CUP ENCLOSED** 

Manufactured by:

Hopkins Agricultural Chemical Co.

Madison, Wisconsin

#### **DIRECTIONS FOR USE**

CORN—To protect germinating seeds and seedlings against seed corn beetles and seed corn maggot, apply 3-4 ½ ounces 1970 Diazinon Seed Protectant per bushel of seed (1 heaping measure equals one ounce) at planting time, either by thoroughly mixing in a box or tub or by dusting it over the seed as it is dumped into the planter box. Thoroughly stir the product with a paddle or lath onto the seed so that each seed is coated with the seed protectant. If temperature is high and soil moisture is low, reduce dosage to one-half, especially for miniseason planting. DO NOT USE BARE HANDS:

1970 Diazinon Seed Protectant should be applied in conjunction with or over an approved fungicidal seed treatment, but do not use on send already treated with an insecticide. Plant within one month of treatment.

Under some circumstances 1970 Diazinon Seed Protectant may affect the rate of seedling. It is necessary for the user to check planting rate frequently to assure himself that the desired planting rate is being obtained.

WARNING: May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of domestic animals. Do not use on livestock, household pets, or humans.

Toxic to fish and wildlife. Do not contaminate any body of water by cleaning of equipment or disposal of wastes and containers. Treated seed exposed on soil surfaces will be hazardous to birds and wildlife.

Treated seed should be labeled as follows:

#### TREATED SEED

#### DO NOT USE FOR FOOD, FEED OR OIL.

When container is empty, immediately wash thoroughly and destroy. NEVER RE-USE.

#### **NOTICE OF WARRANTY**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 2393-199

COEPTED

AND RODENIGUE ACT

CAND RODENIGUE ACT

CAND RODENIGUE ACT

CAND A 2 3 43 - 199

13413-REL