Best Available Copy

DIAZINON SEED PROTECTANT

For the protection of seed and germinating seedlings of corn, peas, soybeans and beans from damage by seed corn maggots and protection of corn from damage by seed corn beetles.

This product contains graphite to assist in accurate planting.

MEASURING CUP ENCLOSED

WARNING: May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of domestic animals. Do not use on livestock, household pets, or humans.

Toxic to fish and wildlife. Do not contaminate any body of water by cleaning of equipment or disposal of wastes and containers. Treated seed exposed on soil surfaces will be hazardous to birds and wildlife.

Treated seed should be labeled as follows:

Treated Seed Do not use for Food, Feed or Oil.

When container is empty, immediately wash thoroughly and destroy. Never Reuse.

NOTICE OF WARRANTY

Selier makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 2393-199



DIAZINON SEED PROTECTANT

For the protection of seed and germinating seedlings of corn, peas, soybeans and beans from damage by seed corn magnots and protection of corn from damage by seed corn beetles.

This product contains graphite to assist in accurate planting.

MEASURING CUP ENCLOSED

DIRECTIONS FOR USE

Corn—To protect germinating seeds and seedlings against seed corn beetles and seed corn maggot, apply 3-4½ ounces Diazinon Seed Protectant per bushel of seed (one level measure equals 1½ ounces) at planting time, either by thoroughly mixing in a box or tub or by dusting it over the seed as it is dumped into the planter box.

Peas, Soybeans and Beans (Lima, Snap, White and Kidney)

SEED CORN MAGGOTS: Mix 1 1/2-3 oz. per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Thoroughly stir the product with a paddle or lath onto the seed so that each seed is coated with the seed protectant. If temperature is high and soil moisture is low, reduce dosage to one-half, especially for mid-season planting. DO NOT USE BARE HANDS.

Diazinon Seed Protectant should be applied in conjunction with or over an approved fungicidal seed treatment, but do not use on seed already treated with an insecticide. Plant within one month of treatment.

Under some circumstances Diazinon Seed Protectant may affect the rate of seeding. It is necessary for the user to check planting rate frequently to assure himself that the desired planting rate is being obtained.

ACCEPTED

JUL7 1970 CEPR

TO THE STATE OF THE PARTY OF TH