

2393-209

07/23/1971

1/1

ACCEPTED

JUL 23 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 2393-209

Hopkins SEVIN®

CARBARYL BAIT

FOR THE CONTROL OF INSECT PESTS NAMED BELOW

Active Ingredient:

Carbaryl (1-naphthyl methylcarbamate) 5% by wt.

Inert Ingredients: 95% by wt.

U. S. D. A. Reg. No. 2393-209

U. S. Pat. Nos. 2,903,478 and 3,009,855

SEVIN is the registered trade mark of Union Carbide Corporation for carbaryl insecticide.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

HARMFUL IF INHALED OR SWALLOWED. Avoid Prolonged Breathing of Dust.

Do Not Take Internally. Avoid Prolonged or Repeated Contact with Skin.

Wear regular long-sleeved work clothing. Change to clean clothing daily.

Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor.

Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.**KEEP PETS OFF TREATED AREAS.**

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions.

If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat in excess of established tolerances.

DIRECTIONS FOR USE

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20 to 40 pounds per acre of Bait will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sowbugs.

| CROP | Dosage Rates | Days Between Final Application and Harvest |
|---|--------------|--|
| Cucumbers, squash, melons, peanuts | 20 | None |
| Alfalfa, peas | 20 - 30 | None |
| Cotton | 20 - 30 | None |
| Beans, carrots, corn, eggplants, okra, peppers, potatoes, tomatoes and sweet corn | 20 | None |
| Asparagus, strawberries | 20 - 40 | 1 Day |
| Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots) | 20 - 40 | 3 Days |
| Sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss chard and turnips (tops) | 20 - 40 | 14 Days |

DISPOSAL OF CONTAINERS

Do Not Reuse Empty Container

Destroy it by burying with waste or burning.

Stay away from smoke or fumes.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHT

Distributed by

50 POUNDS

HOPKINS AGRICULTURAL CHEMICAL CO.
MADISON, WISCONSIN

HACCO 1M 671