2393-

HOPKINS SEVIN® CARBARYL BAIT

FOR THE CONTROL OF INSECT PESTS NAMED ON BACK PANEL

ACTIVE INGREDIENT: Carbaryi (1-Naphthyl methylcarbamate) INERT INGREDIENTS..... TOTAL..... 100% EPA Reg. No. 2393-209 EPA Est. No. 2393-WI-1

®SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS

STATEMENTS OF PRACTICAL TREATMENTS

If on Skin:

Wash thoroughly with soap and water.

If Swallowed: Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger or blunt object. Call a physician or Poison Control Center immediately. Do not induce vomiting or give anything by mouth to an uncon-

cious person.

If in Eves:

Flush with water for at least 15 minutes and get medical attention.

NOTE TO PHYSICIAN: Carbary! is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase-inhibiting drugs.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals CAUTION

HARMFUL IF INHALED OR SWALLOWED. Avoid Breathing of Dust. Do Not Take Internally, Avoid Contact with Skin, Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.
DID CONTAMINATION OF FOOD AND FEED.

φ pets off treated areas.

Environmental Hazards THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. Protective information may be obtained from your Cooperative Agricultural Extension Service. This pesti-? cide is toxic to aquatic organisms. Keep out of lakes, sfreams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.


See Back Panel For Additional Precautionary Statements and Directions For Use

NET WEIGHT

MANUFACTURED BY HOPKINS Agricultural Chemical Co., BOX 7532, MADISON, WISCONSIN 53707

5 POUNDS (2.27 KILOS)





Hopkins

agricultural chemical co.

MADISON, WISCONSIN 53707

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions. If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat in excess of established tolerances.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. DO NOT REUSE EMPTY CONTAINER. PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place. GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning. If burned, stay away from smoke or fumes.

Broadcast application may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20 to 48 pounds per acre of Bait will control cutworms, armyworms, field crickets, darkling beetles, grasshoppers, sowbugs and ants.

CROP_		DOSAGE RATE	Days Between Fina Application and Harv
Cucumber, squash, melons, peanuts	. 20 pounds per acre	e (71/3 ozs. per 1000 sq. ft.)	NONE
Alfalfa, cotton, peas	.30 pounds per acre	(11 ozs. per 1000 sq. ft.)	NONE
Beans, carrots, eggplants, okra, peppers, potatoes, tomatoes	. 40 pounds per acre	(14 ² / ₃ ozs. per 1000 sq. ft.)	NONE
Corn and sweet corn	. 20-40 pounds per a	acre (71/₃ to 14²/₃ ozs. per 1000 s	sq. ft.) NONE
Asparagus, strawberries	. 40 pounds per acre	(142/3 ozs. per 1000 sq. ft.)	1 Day
Broccoli, brussels sprouts, cabbage, cauliflower, head lettuce, garden beets			,
(roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots)	.40 pounds per acre	(14 ² / ₃ ozs. per 1000 sq. ft.)	3 Davs
Sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf		• •	
lettuce, parsley, spinach, swiss chard and turnips (tops)	.40 pounds per acre	(14 ² / ₃ ozs. per 1000 sq. ft.)	14 Days
Tobacco — Field	. 20-10 porinds per a	acre (71/3 to 142/3 ozs. per 1000 s	q ft.) NONE
Bed	1/2-1 pour d per 1.00	00 sq. ft.	NONE
Dandelions	.40-48 porincs per a	acre (142/3 to 171/2 ozs. per 1000 s	q. ft.) 14 Davs
(All Insects listed above except darkling beetles) for the following clops:			•
Apples, apricots, cherries, citrus, grapes, nectarines, peaches, pears, plums,			
annual plants-herbaceous plants and shrubs, trees, woody plants			
(ornamental)	40-49 pounds per a	acre (142/3 to 171/2 ozs. per 1000 s	g. ft.) NONE
NOTICE OF	VARFANTY		

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CODE 1.2.80 1 2 3 4 5 6 7 8 9 (3-80)