

Hopkins


SEVIN brand **carbaryl** insecticide **BAIT**

**FOR THE CONTROL OF CUTWORMS, ARMYWORMS, FIELD CRICKETS,
DARKLING BEETLES, GRASSHOPPERS, SOWBUGS AND ANTS.**

ACCEPTED

DEC 22 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 2393-209



2393-209

12/22/1988

ACTIVE INGREDIENT:
Carbaryl (1-Naphthyl methylcarbamate)5%
INERT INGREDIENTS:95%

TOTAL 100%

EPA REG. NO. 2393-209

EPA EST. NO. 2393-WI-1

* SEVIN is the Registered T.M. of RHONE-POULENC NEDERLAND B.V. for carbaryl insecticide.

**KEEP OUT OF REACH OF CHILDREN
AND DOMESTIC ANIMALS**

CAUTION

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

AVOID CONTAMINATION OF FOOD AND FEED.

Keep pets off treated areas.

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENTS OF PRACTICAL TREATMENT

If on Skin: Wash thoroughly with soap and water.

If Swallowed: Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger or blunt object. Call a physician or Poison Control Center immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If in Eyes: Flush with water for at least 15 minutes and get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase-inhibiting drugs.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except for rice uses. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

Carbaryl is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions. If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat in excess of established tolerances.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral precautions must be given to workers engaged in agricultural hand labor who are expected to be in a treated area or in an area about to be treated with this product. These precautions are: CAUTION: Area treated with carbaryl on (date of application). Do not enter treated area the day of application without protective clothing and gloves. Protective clothing at a minimum should include clean long pants, long sleeved shirt, shoes and socks. For accidental exposure, including skin contact with pellets see Statements of Practical Treatments.

Oral precautions must be given if there is reason to believe that written precautions cannot be understood by workers. When oral precautions are given, precautions shall be given in a language customarily understood by workers.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in cool, dry place in original container. Protect from excessive heat. Once opened, do not store bag for prolonged period. In case of puncture, use overslip bag and securely close. Do not reuse container.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into applicator equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Broadcast application may be made with either ground equipment or aircraft, under conventional, minimal and no-till practices. Treatments may be repeated as necessary. Where a range of rates is provided, use lower rates for light infestation, early instars or when foliage is not too dense; use higher rates for moderate to heavy infestation, advanced instars or when foliage is thick. For best results on lawn and turf, cut lawn, remove clippings and water thoroughly before treatment. Dosages from 20 to 48 pounds of bait per acre will control cutworms, armyworms, field crickets, house crickets, mole crickets, darkling beetles, grasshoppers, and webworms.

CROP	RATE PER ACRE	RATE PER 1,000 SQ. FT.	DAYS BETWEEN FINAL APPLICATION AND HARVEST
Cucumbers, melons, peanuts, pumpkins, squash	20 lbs.	7 1/2 ozs.	0
Birdfoot trefoil (forage and hay), clover, wheat (pasture, forage)	30 lbs.	11 ozs.	0
Lawn and turf—GROUND APPLICATION ONLY	30 lbs.	11 ozs.	0
Pasture (rangeland and wasteland), forage grasses, fence row and non-crop areas			
Beans, carrots, corn (field, sweet, pop), eggplant, okra, peppers, potatoes, sorghum forage, soybeans, sweet potatoes, tobacco beds, tomatoes	20-40 lbs.	7 1/2-14 1/2 ozs.	0
Asparagus, strawberries	40 lbs.	14 1/2 ozs.	1
Broccoli, Brussels sprouts, cabbage, cauliflower, garden beets (roots), head lettuce, horseradish, kohlrabi, parsnips, radishes, rutabagas, salsify (roots), tobacco (field), turnips (roots)	20-40 lbs.	7 1/2-14 1/2 ozs.	3
Alfalfa, peas	30 lbs.	11 ozs.	3
Cotton	30 lbs.	11 ozs.	7
Celery, Chinese cabbage, collards, dandelion, endive (escarole), garden beet tops, Hanover salad, kale, leaf lettuce, mustard greens, parsley, rice, salsify (tops), spinach, sugar beets, Swiss chard, turnip tops.			
* Do not use on rice fields in which crayfish and/or catfish farming is included in the cultural practice.	20-40 lbs.	7 1/2-14 1/2 ozs.	14
Wheat (grain—Fall or Spring)	30 lbs.	11 ozs.	21
Sorghums (milo, grain sorghum, sweet sorghum and hybrids)	20-40 lbs.	7 1/2-14 1/2 ozs.	21 (grain)
Oats, barley and rye (fall treatment)	20-40 lbs.	7 1/2-14 1/2 ozs.	Do not feed treated fodder to dairy animals or animals being finished for slaughter or allow them to graze on crops treated with this product.
Sunflowers	20-40 lbs.	7 1/2-14 1/2 ozs.	60 (crop for oil, food or feed purposes) 0 (crop for seeding purposes only)

(all insects listed above except darkling beetles) for the following crops:

annual plants—herbaceous plants and shrubs, trees, woody plants (ornamental), grapes	40-48 lbs.	14 1/2-17 1/2 ozs.	0
Apples, cherries, peaches, pears, plums	40-48 lbs.	14 1/2-17 1/2 ozs.	1
Nectarines	40-48 lbs.	14 1/2-17 1/2 ozs.	3
Citrus	40-48 lbs.	14 1/2-17 1/2 ozs.	5
Apricots	40-48 lbs.	14 1/2-17 1/2 ozs.	28

NOTE: Do not make more than two applications after grain heads of wheat, alfalfa and clovers emerge from boot. Do not allow animals to graze on treated sunflower crops.

NOTICE OF WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CODE 4.585 2 3 4 5 6 7 8 9 (4/88 USA)

20436

NET WEIGHT



Hopkins
agricultural chemical co.
MADISON, WISCONSIN 53707
U.S.A.

50 POUNDS (22.68 KILOS)

2/2