


BEST DOCUMENT AVAILABLE

Hopkins



UROX 22

WEED KILLER

For General Weed Control on Non-Crop Lands

ACTIVE INGREDIENT	
Mesuron Trichloroacetate [3-(p-chlorophenyl)-1,1-dimethylurea trichloroacetate]	67
INERT INGREDIENTS	
	33
Total	100

TRADE NAME: UROX 22
 REGISTRANT: HOPKINS AGRICULTURAL CHEMICALS CO.
 ADDRESS: 1000 N. 10th St., St. Paul, Minn. 55102

CAUTION

**KEEP OUT OF REACH OF CHILDREN
 HARMFUL IF SWALLOWED
 IRRITATING TO SKIN, EYES, NOSE AND
 THROAT**

Do not breathe dust.

Do not get on skin, in eyes or on clothing.
 In case of contact, immediately wash skin
 with plenty of water; for eyes, wash with
 water for at least 15 minutes and get
 medical attention.

BEST DOCUMENT AVAILABLE

NOTE: Do not use on or near desirable plants or crops.
 Do not use on or near desirable plants or crops where their roots may extend. Equip
 the container with a tight fitting cap to prevent corrosion or contamination of other materials.

DIRECTIONS

NON-CROP LAND

For use on non-crop land such as Rights of way, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas,
 Parking Lots and Under Asphalt for control of annual and perennial grasses and broadleaved weeds. Control
 will be best if applied with a sprayer. The effect, speed and length of time control will last depend on type of soil
 and weather conditions.

RATE: Under ordinary conditions use at the rate of 1.0 to 2.0 pounds per acre. Under conditions where weeds are
 very heavy, increase the rate to 3.0 pounds per acre. For spot treatment use 1.0 to 2.0 pound per 100 square
 feet. Spread the material evenly over the area to be treated.

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over
 the area to be treated. Apply when top growth is dense, but better control will be obtained when weeds
 are young. Apply when weeds are young. Apply when weeds are young.

WHEN TO APPLY: Apply when weeds are young. Apply when weeds are young. Apply when weeds are young.

FLORIDA CITRUS GROVES

For Use in Drainage Ditches For Control of Most Annual and Perennial Grasses and Broadleaved Weeds such
 as Paragrass, Arrowhead, Guinea Grass and Water Primrose.

RATE: 1.0 to 2.0 pounds per acre. (See chart)

	Amount to Apply Per 100 Linear Feet of Ditch				
Width of Ditch	1	2	3	4	5
1 to 2	1	1 to 2	1 to 3	2 to 4	2 to 5

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly
 over the area to be treated. Treat only bottom, not banks.

WHEN TO APPLY: Apply when ditch is not in use but soil is moist. If ditch bottom is not moist, wet it down before
 treating. If it is in use, apply before time of treatment and time of intended use. Run water into the ditch
 for 24 hours before applying. Do not apply when water is in ditch.

NOTE: Do not use in ditches into which tree or other desirable plant roots extend. Do not contaminate drinking
 water. Apply this product to water to be used for livestock.

NOTICE OF WARRANTY

For a detailed description of the use and method of application, weather and crop conditions, mixtures with other
 chemicals, and other factors in the use of this product, see the full label. The Seller, Hopkins Agricultural Chemicals Co.,
 does not warrant the effectiveness of this material in any case where it is not used in strict accordance with
 the label directions.



NET WEIGHT

50 POUNDS