


NET WT. 4 LB.



**Hopkins**

# sevin<sup>®</sup> insect bait

## AN ATTRACTANT INSECTICIDE

For the Control of Cutworms, Armyworms, Crickets, Darkling Ground Beetles, Grasshoppers and Sowbugs.

### ACTIVE INGREDIENT:

Carbaryl (1-naphthyl methylcarbamate) . . . . . 5% by wt.

INERT INGREDIENTS: . . . . . 95% by wt.

U.S. Pat. Nos. 2,903,478 and 3,009,855  
 ©SEVIN is the registered trade mark of Union Carbide Corporation for carbaryl insecticide.

## CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

(See Back Panels for Additional Cautions)  
 EPA Reg. 2393-213

# Hopkins sevin<sup>®</sup> insect bait

## IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions.

If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat in excess of established tolerances.

### DIRECTIONS FOR USE

Apply pellets evenly over vegetables with proper spreading device. Treatments may be repeated as necessary. Dosages of Bait from one-half to one pound per 1000 sq. ft. will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sowbugs.

CROP	Dosage Rates	Days Between Final Application and Harvest
Cucumbers, squash, melons, peanuts, Alfalfa, peas	1/2 lb. per 1000 sq. ft.	None
Cotton	1/2 lb. per 1000 sq. ft.	None
Beans, carrots, corn, eggplants, okra, peppers, potatoes, tomatoes and sweet corn	1 lb. per 1000 sq. ft.	None
Asparagus, strawberries	1 lb. per 1000 sq. ft.	1 Day
Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots)	1 lb. per 1000 sq. ft.	3 Days

CROP (Cont.)	Dosage Rates	Days Between Final Application and Harvest
Sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss Chard and turnips (tops)	1 lb. per 1000 sq. ft.	14 Days

### DISPOSAL OF CONTAINERS

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

### NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

HARMFUL IF INHALED OR SWALLOWED. Avoid Breathing of Dust.

Do Not Take Internally. Avoid Contact with Skin.

Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS. KEEP PETS OFF TREATED AREAS.

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

HOPKINS AGRICULTURAL CHEMICAL CO.  
 Madison, Wisconsin 53701



For home garden