

# Hopkins sevin 50-w insecticide

NET WT. 6 OUNCES

Read entire label. Use strictly in accordance with label cautions, warnings and directions. Keep this pesticide in its original container. Do not put concentrate or dilute into food or drink containers.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

## DIRECTIONS FOR USE

Hopkins Sevin 50W is a wettable spray powder. Measure in level tablespoonfuls. (2 pounds per 100 gallons of water is equal to two level tablespoonfuls per gallon.) Keep mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline products. Do not mix with oil base sprays. This package makes 10 gallons of spray. 20 level tablespoonfuls equal 6 ounces Hopkins Sevin 50W. ORNAMENTALS: Use 2 tablespoonfuls per gallon for Japanese beetles, flea beetles, bagworms, leafhoppers, elm leaf beetles, certain scale insects, box elder bugs and tent caterpillars. Apply when pests appear. Repeat as needed.

APPLES AND PEARS: Use 2 tablespoonfuls per gallon for codling moth, apple maggot, plum curculio, fruit tree leaf rollers, Japanese beetles, apple aphids and San Jose scale. Do not apply before second cover spray to avoid fruit thinning. Application of Carbaryl within 30 days after full bloom may also provide apple thinning. To avoid this, delay use until 30 days after bloom. May cause foliar injury if used before second cover on York and McIntosh apples. Apply at 10 day intervals in July for apple maggot. May be used up to 1 day before picking.

STRAWBERRY: Use 2 tablespoonfuls per gallon up to 1 day before picking for leaf roller and meadow spittlebug control. Makes 3 to 5 post bloom summer sprays as needed.

BEANS: (Green, lima, snap, cowpeas and black-eyed peas): Use 1 tablespoonful per gallon for Mexican bean beetle. Use 2 - 3 tablespoonfuls per gallon for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, armyworms, cutworms, corn earworm, stink bugs and tarnished plant bug.

CORN; Sweet: Use 2 tablespoonfuls per gallon for corn earworm, corn borer, fall army worms and grasshoppers. Apply up to day of harvest. Apply to silks as soon as they appear. Make 3 to 4 applications at 2 - 3 day intervals as infestation warrants. Re-apply immediately after rain. Repeat

at 2 to 3 day intervals for earworms or until silks are dry. Apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. Do not feed treated fodder to animals. CUCUMBER, MELONS AND SQUASH: Use 2 tablespoonfuls per gallon. Apply as full wetting spray for squash bugs, melon worms, cucumber beetles and pickleworms.

POTATO, TOMATO, PEPPER, EGGPLANT: Use 2 tablespoonfuls per gallon for Colorado potato beetle, flea beetles and leafhoppers. Use 2 - 4 tablespoonfuls per gallon for tomato fruitworm, fall armyworms, stink bugs and lace bugs.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: Use 1 to 2 tablespoonfuls for flea beetles and harlequin bug. Use 2 to 4 tablespoonfuls for armyworms, cabbage worms and loopers and corn earworm. Where cabbage looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small cabbage loopers after edible portions begin to form, use Sevin in a 5 to 7 day schedule. Do not apply within 3 days of harvest of these crops.

## CAUTION

### KEEP OUT OF REACH OF CHILDREN AND ANIMALS

HARMFUL IF INHALED OR SWALLOWED. Avoid Breathing of Dust or Spray. Do Not Take Internally. Avoid Contact with Skin. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.


NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

## NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NOTE: This product sold by weight and package is full when packed, but due to fluffiness of materials it is inclined to settle when shipped.



6-8-72  
2393-222C

# Hopkins sevin<sup>®</sup> 50-w insecticide

A WETTABLE POWDER FOR CONTROL OF INSECT PESTS

## ACTIVE INGREDIENT:

Carbaryl (1-naphthyl methylcarbamate) ..... 50% by wt.

INERT INGREDIENTS: ..... 50% by wt.

EPA Reg. No. 2393-220

®SEVIN is the registered trade mark of Union Carbide Corporation for carbaryl insecticide.

## CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

HOPKINS AGRICULTURAL CHEMICAL CO.  
Madison, Wisconsin 53701



For home garden