

DIAZINON-CAPTAN SEED PROTECTANT

Diazinon-Captan Seed Protectant gives temporary protection of germinating seed and seedlings of corn, peas, soybeans and beans (lima, snap, white and kidney) from injury by seed corn maggots and seed corn beetles and gives temporary protection to seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

This product contains graphite to assist in accurate planting.

CORN: Use 6 oz. (3 level cups) per bushel corn seed. Apply to seed as planter box is being filled. Stir with paddle or lath so that each seed is coated with seed protectant. If temperature is high and soil moisture is low reduce dosage to one-half especially for mid-season planting. **DO NOT USE BARE HANDS.** Excess product flows with the seed into the furrow giving additional protection against soil-borne diseases.

PEAS, SOYBEANS AND BEANS (lima, snap, white and kidney): Mix 3 oz. (1½ level cups) per bushel of seed in the planter box as a seed treatment prior to planting. Follow mixing instructions as above.

Diazinon-Captan Seed Protectant may be applied over an approved fungicidal seed treatment.

Under some circumstances Diazinon-Captan Seed Protectant may affect the rate of seeding. It is necessary for the user to check planting rate frequently to assure himself that the desired planting rate is being obtained.

MEASURING CUP ENCLOSED.

Manufactured by

HOPKINS AGRICULTURAL CHEMICAL CO.
Madison, Wisconsin 53701

HACCO 13M-1172

Hopkins

MOSQUITO LARVICIDE

FOR CONTROL OF MOSQUITO AND MIDGE LARVAE

Contains 1% ABATE[®] Insecticide


ACTIVE INGREDIENT:
O,O,O',O' tetramethyl O,O' thiodiphenylene p
INERT INGREDIENTS:

EPA REG. NO. 2393-221
* See label for full list of ingredients

CAUTION

KEEP OUT OF REACH OF CHILDREN

MAY BE HARMFUL IF SWALLOWED
Avoid breathing of dust
Avoid contact with skin
Wash thoroughly after handling
Do not contaminate food or feed



Comments DIRECTIONS FOR USE

Do not use this product for any uses other than those specified on

For control of mosquito and midge larvae, apply Hopkins Mosquito

Areas of Treatment

Standing waters, shallow ponds, lakes, woodland pools

Tidal waters, marshes, swamps and waters high in organic content

May be applied as a spot treatment to catch basins and similar

Apply uniformly and repeat applications as necessary.

Do not use on crops used for food, forage or pasture.

This product is toxic to birds. Keep out of lakes, streams, or ponds
Application rates recommended on this label. Do not apply where the
contaminate water by cleaning of equipment or disposal of wastes.
this label.

Destroy empty bags by burning, away from buildings and fuel tanks

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the
on the label. Buyer assumes all risk of use and handling of this
is confined to label restrictions.

NET WEIGHT

Hopkins Agricultural Chemicals
MADISON, WISCONSIN 537