

Hopkins

10% SEVIN[®] DUST

For the Control of Certain Vegetable Crop Insect Pests


ACTIVE INGREDIENT:

Carbaryl (1-naphthyl methylcarbamate) 10%

INERT INGREDIENTS: 90%

TOTAL 100%

[®]SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.
EPA Reg. No. 2393-225



CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

TAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

To control the insects on the crops specified, apply Hopkins 10% Carbaryl Dust at the amounts stated below. Apply in a manner to give uniform coverage. Make first application when insects or their damage appear. Repeat as specified or as necessary.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: Controls flea beetles and harlequin bug. Use at the rate of 6 - 10 lbs. per acre; for cabbage worms and loopers and corn earworm—use 10 - 20 lbs. per acre. Do not apply within 3 days of harvest. Where cabbage looper is the principal problem use a specifically recommended alternative pesticide.

BEANS — Green, Snap or Lima: Controls Mexican bean beetle, bean leaf beetles, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, armyworms, stink bugs and tarnished plant bug — use at the rate of 10 - 15 lbs. per acre.

CORN — Sweet corn: Controls corn earworm, European corn borer, Japanese beetle and sap beetles. To control foliar feeding insects, treat entire plant repeating as necessary. To control insects feeding on ears and silks treat at 2 - 3 day intervals, beginning when silks appear and continue until silks are dry — use at the rate of 10 - 20 lbs. per acre.

POTATO, TOMATO, EGGPLANT, PEPPER: Control Colorado potato beetle, flea beetles, leaf hoppers -- use at the rate of 6 - 10 lbs per acre; for tomato fruitworm, tomato hornworm and stink bugs — use 10 - 20 lbs. per acre.

CUCUMBER, MELONS, PUMPKIN AND SQUASH: Controls pickleworm and melonworm — use at the rate of 6 - 10 lbs. per acre. **NOTE.** Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not overtrear.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHT

50 POUNDS

Hopkins Agricultural Chemical Co.

MADISON, WISCONSIN 53701

HACCO 2M 973