

Hopkins

TURF INSECTICIDE

FOR THE CONTROL OF TURF INSECTS

ACTIVE INGREDIENT

Carbaryl (1-naphthyl-methyl-carbamate) 10% by wt

INERT INGREDIENTS

90% by wt

100% 100

EPA Reg. No. 2443-PP

*SEVIN is the registered trademark of the manufacturer of Carbaryl insecticide.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED - Avoid Breathing of Dust or Spray. Do Not Take Internally. Avoid contact with eyes. Wear regular clothing and shoes. Wash hands thoroughly after handling. Wash thoroughly before eating or drinking.

NOTE FOR PHYSICIAN - Carbaryl is a carbamate. It is a non-steroidal anti-cholinergic.

AVOID CONTAMINATION OF FOODS, FEEDS, FISHES, APPLIES, STREAMS AND PONDS

Do not apply to areas where food, feed, fish, or other animals are present. Do not apply to areas where water is used for drinking or irrigation. Do not apply to areas where water is used for irrigation.

Protective clothing should be worn when handling. Wash thoroughly after handling.

Do not use in areas where children or pets are present.

Readent product label for complete directions.

DIRECTIONS FOR USE

TURF INSECTS - Treat the surface of the turf with the carbaryl insecticide.

A
4/1 - 1777
1977

Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate to strong cholinesterase inhibitor. Atropine is an antidote.

AVOID CONTAMINATION OF FOOD, FEED, WATER, SPRINGS, STREAMS AND PONDS

For protection of crops, feed, water, springs, streams, ponds, and other bodies of water, the area should be sprayed during periods of low water level and before runoff. Maintain a safe distance until 1 week after treatment.

Protective information may be obtained from your local State Agricultural Extension Service.

Do not reuse empty container. Destroy it by burying or incineration. Stay away from smoke or fumes.

Read entire label. Use in strict accordance with label directions and caution.

DIRECTIONS FOR USE

TURF INSECTS: Treat the turf evenly and thoroughly, at the rate of two pounds of Hopkins Turf Insecticide for each 1,000 square feet to control ants, bluegrass billbug, chinch bugs, cutworms, earwigs, fall armyworm, leafhoppers, millipedes, mosquitoes, and sod webworm (lawn moths).

SPECIAL INSTRUCTIONS: For best results, mow lawn, remove clippings, and water thoroughly before treatment. Apply with a spreader for uniform coverage. Work grass areas with a rotary water spray after treatment. Repeat treatment in 3-4 weeks later if necessary. The two pounds of Hopkins Turf Insecticide for each 1,000 square feet of treated area lawn is a very effective control against ticks. Work grass areas with a rotary water spray after treatment. Repeat treatment in 3-4 weeks later if necessary.

OTHER USES:

For control of ticks on lawns, apply at the rate of two pounds of Hopkins Turf Insecticide for each 1,000 square feet of treated area. Work grass areas with a rotary water spray after treatment. Repeat treatment in 3-4 weeks later if necessary.

NET WEIGHT 40

5 POUNDS

A
Sept 12, 1972
1373 578

Hopkins Agricultural Chemical Co.
MADISON, WISCONSIN 53701