


# Hopkins

## CAPTAN-STREPTOMYCIN 7.5-.01 POTATO SEED PIECE PROTECTANT



### ACTIVE INGREDIENTS:

Captan (N trichloromethylthio 4-cyclohexene 1,2-dicarboximide) . . . . . 7.50%  
Streptomycin\* . . . . . .01%

### INERT INGREDIENTS:

. . . . . 92.49%  
TOTAL . . . . . 100.00%

\*Streptomycin from streptomycin sulfate (agricultural grade). Contains 100 ppm. streptomycin

EPA Reg. No. 2393-236

### CAUTION

KEEP OUT OF REACH OF CHILDREN

AVOID CONTAMINATION OF FEED AND FOODSTUFFS. AVOID INHALATION OF DUST. AVOID CONTACT WITH SKIN. HARMFUL IF TAKEN INTERNALLY. STORE IN A COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Dispose of excess treated seeds or wastes by burial.

Store in Cool Dry Place - Protect from Excessive Heat

Do not use treated seed for food or feed purposes

Burn bag immediately when empty. Stay out of smoke

### DIRECTIONS FOR USE

This material is especially formulated for potato seed piece treatment

**POTATOES (SEED PIECES):** To control Seed Rot and Damp off, Verticillium, Wilt, Brown Eye Disease, Dust at the rate of 10 ounces per bushel cut seed pieces (16 ounces or 1 lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment to sprayer. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 or 3 days in open crates before bagging.

### NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

NET WEIGHT

50 POUNDS

Hopkins Agricultural Chemical Co.  
Madison, Wisconsin 53701