

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Do not induce vomiting without the advice of a physician — contains petroleum solvent, an aspiration hazard. Do Not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Remove contaminated clothing and wash affected skin with soap and water.

If in eyes: Flush with water for at least 15 minutes. Get medical attention

If inhaled: Move to fresh air and assist respiration if needed.

Note to Physician: Malathion is a cholinesterase inhibitor. Atropine is antidotal.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Do not contaminate food or feed products. Wash thoroughly with soap and water after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Service.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

NOTICE OF WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

GENERAL DIRECTIONS

This product may be used in conventional hydraulic sprayers, low-volume ground applicators and airplane sprayers. Mix this product with water by agitation. Agitate the mixture while spraying. If allowed to stand, re-agitate before use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALFALFA — To control Alfalfa Weevil Larvae, use 2 to 3 quarts this product per acre in 15 to 25 gallons of water. Repeat applications as necessary. Do not use within 7 days of harvest.

Continued on right panel



Hopkins


Agricultural Chemical Co.

Box 111 • Madison, Wisconsin 53701

CODE 4 27 81 1 2 3 4 5 6 7 8 9 10 81

Hopkins

ALFALFA AND CLOVER SPRAY 2-2 E.C.



ACTIVE INGREDIENTS:

Methoxychlor* 23.80%

Malathion (O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate) 23.82%

Xylene 44.63%

INERT INGREDIENTS: 7.75%

Total 100.00%

1 Gallon Contains 2 Pounds of Methoxychlor and 2 Pounds of Malathion
*23.80% methoxychlor, technical is equivalent to 20.825%
2,2-Bis (p-methoxyphenyl)-1,1, 1-trichloroethane and 2.975%
other isomers and reaction products.

*Registered trademark Hopkins Agricultural Chemical Co.
EPA Reg. No. 2393-237

EPA Est. No. 2393-IL-3*, 2393-WI-1*,
523-IL-1*, 2393-WI-2*, 33509-MO-1*
Superscript refers to first letter of lot number

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE SIDE PANEL FOR USE INSTRUCTIONS
SEE SIDE PANEL FOR PRECAUTIONS AND NOTICE OF WARRANTY

Net Contents

5 Gallons

DIRECTIONS (Continued)

ALFALFA, CLOVER — Alfalfa Weevil Larvae, Spittlebug, Pea Aphid, Leafhoppers, Spider Mites, Grasshoppers, Spotted Alfalfa Aphid, Lygus Bug, Armyworm, Webworm, Alfalfa Caterpillar use 2-3 quarts per acre. Do not apply during full bloom. Do not apply until temperature is 60°F or above. Amount of water to use depends on how thick a stand. Use enough water for complete coverage. Do not use within 7 days of harvest.

CORN (Field, Pop or Sweet), (Forage or Grain) — Aphids, Sap Beetles, Thrips, Young Grasshoppers, Corn Rootworm (Adults), Japanese Beetle, Fall Armyworms, Flea Beetles, Rose Chafer, Cereal Leaf Beetle use 2 quarts per acre. Apply on silks and foliage as needed and repeat if necessary. Do not use within 7 days of harvest.

RED TOP, TIMOTHY, ORCHARD GRASS, FESCUE, CANARY GRASS — Cereal Leaf Beetle use 2 quarts per acre. Repeat as necessary. Do not use within 7 days of harvest.

ASPARAGUS — Asparagus Beetles, Thrips use 2 quarts per acre. Do not use within 3 days of harvest.

BEANS — Mexican Bean Beetle, Leafhoppers, Spider Mites, Japanese Beetle, Aphids, Cucumber Beetles, Corn Earworm, Flea Beetles, Bean Leaf Beetles, Fall Armyworms, Rose Chafers use 4-5 pints per acre. Do not use within 3 days of harvest.

BEETS — Aphids, Blister Beetles, Flea Beetles use 4-5 pints per acre. Do not use within 7 days of harvest. If tops are to be used as food or feed, do not apply within 14 days of harvest.

BROCCOLI*, BRUSSELS SPROUTS*, KALE*, CABBAGE, CAULIFLOWER**, CARROTS**, KOHLRABI**, TURNIPS**** — Aphids, Imported Cabbage Worms, Cabbage Loopers, Leafhoppers, Flea Beetles, Fall Armyworms use 4-5 pints per acre. Do not use within 7 days** - 14 days* of harvest. Do not use within 14 days of harvest on carrots and turnips if tops are to be used as food or feed.

CANTALOUPE, CUCUMBER, MELONS, SQUASH (Summer & Winter), PUMPKINS* — Aphids, Spider Mites, Cucumber Beetles, Flea Beetles, Leafhoppers, Pickleworms, use 4-5 pints per acre. Do not use within 1 day of harvest. *Pumpkins — Do not use within 3 days of harvest.

EGGPLANT, PEPPERS — Aphids, Spider Mites, Flea Beetles, Leafhoppers, Rose Chafers use 4-5 pints per acre. Do not use within 3 days of harvest.

LETTUCE, SPINACH — Cabbage Looper, Leafhoppers, Alfalfa Looper use 4-5 pints per acre. Aphids use 5 pints per acre minimum. Do not use within 14 days of harvest.

PEAS — Pea Aphids, Pea Weevil, Alfalfa Looper use 4-5 pints per acre. Do not use within 7 days of harvest.

POTATOES — Aphids, Leafhoppers, False Chinch Bugs, Mealybugs, Blister Beetles, Flea Beetles, Colorado Potato Beetles, Cucumber Beetles use 4-5 pints per acre.

TOMATOES — Aphids, Spider Mites, Blister Beetles, Fall Armyworms, Japanese Beetles, Tomato Fruitworm use 4-5 pints per acre. Do not use within 1 day of harvest.

STRAWBERRIES — Aphids, Spider Mites, Lygus Bugs, Spittlebugs, Field Crickets, Potato Leafhopper, Strawberry Leafroller, Whiteflies, Flea Beetles, Omnivorous Leaf Tier, Strawberry Weevil use 4-5 pints per acre. Do not use within 3 days of harvest.

BLUEBERRIES — Japanese Beetle, Cherry Fruitworm, Cranberry Fruitworm, Leafhoppers use 4-5 pints per acre. Apply when largest berries of earliest varieties are 1/2 inch in diameter and 7-10 days later. Do not use within 14 days of harvest.

BLACKBERRIES, BOYSENBERRIES*, DEWBERRIES, LOGANBERRIES, RASPBERRIES — Mites, Thrips, Leafhoppers, Japanese Beetle, Spittlebug, Rose Chafer, Strawberry Weevil, Flea Beetles use 4-5 pints per acre. Do not use within 3 days of harvest. *BOYSENBERRIES — Do not use within 14 days of harvest.

GRAPES — Leafhoppers, Spider Mites, Grape Berry Moth, Mealybugs, Japanese Beetle, Rose Chafer, Flea Beetles, Western Grape Skeletonizer use 4-5 pints per acre. Do not use within 14 days of harvest. Injury may occur on Ribier, Italia, Cardinal & America varieties if applied after the cluster stage.

APPLES, PEARS — Aphids (Woolly apple, rosier apple & black cherry), Scales (forbes & San Jose), Bud Moth, Tent Caterpillar, Mealybug, Mites (European red clover, Williamette and two spotted), Codling Moth, Plum Curculio, Pear Psylla, Japanese Beetle, Leafhoppers, Pear Sugs use 2 quarts in 100 gallons of water. For Pear Fruit Tree Leafroller and Red Banded Leafroller use 5 pints in 100 gallons of water. Give full coverage. It may cause injury to foliage on some varieties of apples and Bosc pears. Do not use within 7 days of harvest. Apply at petal fall and every 10 to 14 days thereafter as long as local conditions warrant.

APRICOTS, PEACHES, NECTARINES — Codling Moth, Plum Curculio, Japanese Beetle, Orange Tortrix, Scales (European Lecanium, Soft Brown, Terrapin & Cottony Pear), Aphids (Green Peach, Black Cherry, Black Peach & Rusty Plum), Mites use 2 quarts in 100 gallons of water. Give full coverage application. Do not use within 21 days of harvest. Apply at petal fall and every 10 to 14 days thereafter as long as local conditions warrant.

CHERRY — Plum Curculio, Cherry Fruit Flies, Cherry Fruit Worms, Black Cherry Aphids, Bud Moth, Fruit Tree Leafrollers, Japanese Beetles use 2-2 1/2 quarts in 100 gallons of water. Give full coverage application. May injure foliage of some varieties. Do not use within 7 days of harvest. Apply at petal fall and every 10 to 14 days thereafter as long as local conditions warrant.

PLUMS, PRUNES — Plum Curculio, Japanese Beetle, Mealy Plum Aphids, Bud Moth use 1 1/2 quarts in 100 gallons of water. Full coverage application. Do not use within 7 days of harvest. Apply at petal fall and every 10 to 14 days thereafter as long as local conditions warrant.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

STORAGE: Do not store near heat or open flame. Do not store near food or feed.

PESTICIDE DISPOSAL: Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse for equivalent and offer for recycling or reuse if allowed. Or dispose of in a sanitary landfill or by other approved State and local procedures.