


Hopkins FLY BAIT GRITS

Attracts and Kills Houseflies

NOT FOR STORAGE IN OR AROUND THE HOME
RESIDENTIAL USE RESTRICTED TO COMMERCIAL APPLICATORS



ACTIVE INGREDIENT:

Bomy® (Dimethyl 3-hydroxyglutaconate dimethyl phosphate)*	1.00%
INERT INGREDIENTS:	99.00%
TOTAL	100.00%

*Registered trademark of Hopkins Agricultural Chemical Co.
*U.S. Patent No. 2,891,887
EPA Reg. No. 2393-343 AA

EPA Est. No. 2393-WI-1

KEEP CONTAINER CLOSED — STORE IN COOL, DRY PLACE

WARNING

KEEP OUT OF REACH OF CHILDREN

May be harmful or fatal if swallowed, inhaled or absorbed through skin. DO NOT get on skin, in eyes, or on clothing. Wash thoroughly with soap and water after using. Keep out of reach of children.

Should illness occur, get prompt medical aid.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

IF SWALLOWED - Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

Do not contaminate feed and foodstuffs. Do not apply where poultry or animals, especially young calves, can pick it up or lick it. Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Use only in areas inaccessible to food producing animals.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Use this product only as specified on this label.

Do not re-use empty container. Destroy container by perforating, and bury or discard in a safe place.

DIRECTIONS FOR USE

Scatter Hopkins FLY BAIT GRITS daily, or as needed, to reduce fly population. Distribute bait from container, or with jar or can to avoid handling bait. Keep FLY BAIT GRITS in original container, tightly closed, and store in cool place.

LOADING SHEDS, LIVESTOCK PENS, PIG PARLORS, STABLES, BROILER HOUSES: Use only around outside areas. Scatter FLY BAIT GRITS at the rate of ¼ lb. per 1,000 sq. ft. on ground areas where flies congregate. Ideal baiting sites are around doorways, along fence lines, soiled areas, or in alleyways of feedlots.

CAGED POULTRY AND TURKEYS ON WIRE: Scatter FLY BAIT GRITS on walkways at the rate of ¼ lb. per 500 sq. ft. to aid in controlling houseflies. Also scatter outdoors around doorways.

DOG KENNELS AND FURBEARING ANIMAL UNITS: Scatter FLY BAIT GRITS around outside areas of dog pens, runways and other surfaces out of reach of animals. For cage units, scatter on walkways at the rate of ¼ lb. per 500 sq. ft.

FOOD, POULTRY AND MEAT PROCESSING PLANTS: Scatter FLY BAIT GRITS at the rate of ¼ lb. per 1,000 sq. ft. in inedible departments only. FLY BAIT GRITS may be used outside livestock pens or loading and unloading docks.

NET WEIGHT



MANUFACTURED BY

Hopkins
agricultural chemical co.

50 POUNDS

CODE 9 12 75 1 2 3 4 5 6 7 8 9 (78)

BOX 7532 • MADISON WISCONSIN 53707