


Hopkins

POTATO-TOMATO DUST

Controls Certain Insects and Diseases on Potatoes and Tomatoes
and other Vegetables

Contains Sevin[®] and Dithane[®] Materials



ACTIVE INGREDIENTS:

Carbaryl (1-methyl-1-methyl-N-methylcarbamate)	5.0
Neo-chlorogenic acid (a product of zinc ion and manganese chloride)	4.0
Disulfithiocarbamate in which the ingredients are:	
Manganese	0.8
Zinc	0.1
Ethylene disulfithiocarbamate ion	3.1

INERT INGREDIENTS:

.....	91.0
TOTAL	100.0

[®] SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

[®] DITHANE is the registered trademark of John and Haas Co.

EPA Reg. No. 2393-244-AA

EPA Est. No. 2393-WI-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. May cause irritation of nose, throat, eyes and skin. Avoid breathing of dust. Do not take internally. In case of skin contact wash with water. For eyes, wash copiously with water and get medical attention. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change clean clothing daily. Avoid contamination of food, feed, water supplies, streams and ponds.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversibly, cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension or Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning far away from smoke or fumes.

DIRECTIONS FOR USE

To control the insects and diseases on the vegetables specified apply Hopkins Potato Tomato Dust at the amounts stated below. Apply in a manner to give uniform coverage using any suitable dust applicator. Make first application when insects or their damage, or signs of disease appear. Repeat as specified or as necessary. Apply this product only

take internally. In case of skin contact wash with water. For eyes, wash copiously with water and get medical attention. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid contamination of food, feed, water supplies, streams and ponds.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversibly, cholinesterase inhibitor. Atropine is antidotal.

It is predicted to be only toxic to bees exposed to direct treatment or residues on blossoms. Protective information may be obtained from the Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

To control the insects and diseases on the vegetables specified apply Hopkins Potato Tomato Dust at the amounts stated below. Apply in a manner to give uniform coverage using any suitable dust applicator. Make first application when insects or their damage, or signs of disease appear. Repeat as specified or as necessary. Apply this product only

POTATO, TOMATO: To control Colorado potato beetle, flea beetles, leafhoppers, use at the rate of 12-20 lbs. per acre. For tomato fruitworm, tomato hornworm and stink bugs, early and late blight on potatoes, early and late blight, gray leaf mold, gray leaf spot and anthracnose on tomatoes, use at the rate of 20-40 lbs. per acre. **TOMATOES:** Do not apply within 5 days of harvest.

SWEET CORN: Controls corn earworm, European corn borer, Japanese beetle, sap beetles. To control foliar feeding insects, treat entire plant repeating as necessary. To control insects feeding on ears and silks treat at 2-3 day intervals, beginning when silks appear and continue until silks are dry. In small gardens for the control of European corn borer apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge.

To control Helminthosporium leaf blight start application when disease first appears and repeat at 4 to 7 day intervals. Do not apply within 7 days of harvest.

For all purposes use at the rate of 20-40 lbs. per acre.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

CUCUMBER, MELONS, AND SUMMER SQUASH: Controls pickleworm and melonworm, use at the rate of 12-20 lbs. per acre. For cucumber beetles, flea beetles, leafhoppers and squash bugs, use 20 lbs. per acre. Start applications when plants are in the two leaf stage and repeat at 5 to 7 day intervals. Direct dust for thorough coverage of both upper and lower leaf surfaces. Do not apply within 5 days of harvest. **NOTE:** Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not over-treat.

NOTICE OF WARRANTY

Manufacturer makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling of this material is contrary to label instructions.

NET WEIGHT

HOPKINS AGRICULTURAL CHEMICAL CO.

Madison, Wisconsin 53701

50 POUNDS