



Hopkins

MALATHION ULV[®] CONCENTRATE INSECTICIDE

ACTIVE INGREDIENT:

Malathion** 95.0%

INERT INGREDIENTS: 5.0%

TOTAL 100.0%

**O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate
(One gallon contains 9.7 pounds of malathion)

©Trademark of American Cyanamid Company
EPA Reg. No. 2393-250

ACCEPTED
MAY 30 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 2393-250

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL BY SWALLOWING, INHALATION OR SKIN CONTACT

Avoid Breathing Spray Mist.

Avoid Contact With Skin.

Wash Thoroughly After Handling.

Change Contaminated Clothing.

Do Not Contaminate Food or Feed Products.

Highly toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension service.

Apply this product only as specified on this label.

NET CONTENTS

GALLON

Repacked by

HACCO 1M 573

HOPKINS AGRICULTURAL CHEMICAL CO.
Madison, Wisconsin 53701

CROP	PESTS CONTROLLED	FLUID OUNCES PER ACRE	INTERVAL BETWEEN LAST APPLICATION AND HARVEST
Beans (lima, green, snap, Navy, red kidney, wax, dry, blackeye)	Mexican Bean Beetle	8	1 day
	Leafhoppers		
	Green Cloverworm		
	Japanese Beetle		
	Lygus Bugs		
Blueberries	Blueberry Maggot	10	0 day
Nonagricultural Lands	Beet Leafhopper (on wild host plants)	8	0 day
Beef Cattle — Feed lots and Holding Pens	Adult Flies and Mosquitoes	6 - 8	0 day

Alfalfa, Clover, Pasture and Range Grass, Grass and Grass Hay, Grain Crops, Beans, Rice, Tomatoes and Nonagricultural Lands (wastelands, roadsides, soil bank lands): Adult mosquitoes and Flies — Apply Malathion ULV Concentrate at the rate of 2 to 4 fluid ounces for control of adult mosquitoes and at 6 to 8 fluid ounces per acre for control of adult flies and mosquitoes. Repeat applications as necessary. On alfalfa, clover, pasture and range grass, grass and grass hay, may be applied on day of harvest or grazing. Do not apply to alfalfa and clover in bloom. Do not use on seed alfalfa. On grain crops, make no application within 7 days of harvest or forage use; on corn, within 5 days of harvest or forage use; on rice, within 7 days of harvest; on beans and tomatoes, within 1 day of harvest.

FOREST INSECTS

Apply with aircraft equipped for ultra low volume application. Make application when air is calm and temperature is below 68°F. Do not allow spray to contact ferns, hickory and maples as injury may result. Do not spray on elms under extreme heat, drought and disease conditions.

TREE	PESTS CONTROLLED	FLUID OUNCES PER ACRE	DIRECTIONS
Douglas Fir True Fir Spruce	Spruce Budworm	13	Apply when highest percentage of larvae are in the fifth instar.
	Hemlock	8	Apply when most larvae are in the third and fourth instar.
	Pines	European Pine Sawfly	10
Larch	Larch Casebearer	8	Apply in spring as soon as larvae break hibernation and begin feeding on new foliage.

Before using Malathion ULV Concentrate for the preparation of malathion insecticide, manufacturers should consult American Cyanamid Company for manufacturing and safe handling instructions.

DISPOSAL OF PAILS: Drain pail completely. Add 1/2 gallon of water, 1/4 cup of detergent and 1/4 pound of lye. Tighten closure. Rotate pail to wet all surfaces and let stand for at least 15 minutes. Drain completely and rinse several times with water. Bury rinse solution 18 inches deep. Tighten closure. Punch holes in top and bottom of container, crush and bury.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

2393-250

05/30/1973

1/1