

**HOPKINS 3% THIODAN - 7% COPPER DUST**

**CAUTION**

**HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.** Avoid contact with skin, eyes and clothing. Avoid inhalation of dust. Wash thoroughly after handling. Do not take internally. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Avoid contamination of feed and foodstuffs. Do not feed treated forage or plant material to dairy animals or to animals being finished for slaughter. The use of this material in a manner or a time other than in accordance with directions may produce plant injury.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Birds and other wildlife feeding on the treated area may be harmed. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. **DO NOT REUSE EMPTY CONTAINER.** Destroy it by burying with waste or burning. Stay away from smoke or fumes. 2398 249

**DIRECTIONS FOR USE**

This dust is formulated to control certain insects and diseases on potatoes and tomatoes.

In general application should be made at 7 to 14 day intervals, depending on weather conditions and the severity of insect or disease problem.

Local weather and other conditions may affect the use of this material.

**POTATOES:** For control of Aphids, Colorado Potato Beetle, Green Stink Bug, Leaf-footed Bug, Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm, Southern Armyworm, Whitefly, European Corn Borer, Potato Psyllid, False Chinch Bug, Three-lined Potato Beetle, Early and Late Blight—Apply at the rate of ¼ pound per 1,000 square feet at 7 to 14 day intervals, depending on weather conditions and insect and disease incidence. Under conditions which favor Blight development, more frequent applications are required than during dry, hot weather. Begin dusting soon after potato plants show above ground; continue throughout the growing season. Thorough coverage is important. Do not plant root crops other than carrots, potatoes and sweet potatoes as follow-up crops.

**TOMATOES:** For control of Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Hornworm, Tomato Fruitworm, Yellow-striped Armyworm, Cabbage Looper, Stink Bugs, Whitefly, Early Blight, Late Blight and Septoria Leaf Spot—Apply at the rate of ¼ pound per 1,000 square feet. Begin application at first sign of insect infestation or disease infection. Repeat at 7 to 14 day intervals or as necessary to keep problems under control. Do not apply within 1 day of harvest.

**NOTICE OF WARRANTY**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**HOPKINS AGRICULTURAL CHEMICAL CO.**  
Madison, Wisconsin 53701

ACCEPTED  
NOV 5 1973  
FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
BY INSPECTOR  
TO ATTACHED COMMENTS