

Hopkins

25% CAPTAN SEED PROTECTANT

FOR PLANTER OR DRILL BOX APPLICATION

Use on Corn, Peanuts, Soybeans and Wheat
Contains 15% Graphite (non-oily lubricant)

1 Ounce Measuring Cup Enclosed

ACTIVE INGREDIENT:

Captan (N- [(trichloromethyl) thio] -4-cyclohexene 1, 2-dicarboximide) 25.0%

INERT INGREDIENTS: 75.0%

TOTAL 100.0%

EPA Reg. No. 2393-259AA

EPA Est. No. 2393-WI-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Avoid inhaling dust - wear respirator. Avoid contact with skin, eyes or clothing. Wash with soap and water after handling. Do not contaminate feed or food.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Dispose of excess treated seeds by burying with wastes.

Container should be disposed by crushing and burying with wastes.

PROTECT FROM EXCESSIVE HEAT - STORE IN A COOL DRY PLACE.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS.

PRODUCT INFORMATION

Hopkins 25% Captan is especially formulated for the treatment of seed in the planter or drill box. It protects sound and/or mechanically injured seed against seed and soil-borne diseases such as seedling blights, damping-off and seed decay organisms at planting time.

Designed specifically for seed protection at planting time, Hopkins 25% Captan Seed Protectant requires no additional seed handling to have seed treated, this lowers the percentage of cracked or "split" seed planted. At recommended dosage rates, this product assures adequate seed coverage and allows any excess to flow down with the seed into the seed furrow thus affording maximum protection. Can be used on previously treated seed.

A non-oily lubricant has also been added to help prevent cracking and splitting of the seed in the planter box so they will flow easily and evenly.

PLANT SAFETY: At the recommended dosage 25% Captan is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated giving a superior result as compared to results from other seed fungicides.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Do not use treated seed for feed, food or oil purposes.

RECOMMENDATIONS FOR USE

Fill the planter box approximately one-half full of seed and add half the recommended dosage. Stir with a stick or paddle until uniform coverage is observed. Add the remaining seed and chemical and mix thoroughly as above.

CORN (Field) - Use at the rate of 4 ounces per bushel or 7 ounces per 100 pounds of seed as a planter box treatment.

PEANUTS (Shelled) - For control of *Aspergillus niger* and other seed and soil-borne diseases, use 2 to 4 ounces per bushel or 4 to 6 ounces per 100 pounds of seed as a planter box treatment.

SOYBEANS - Use at the rate of 4 ounces per bushel or 7 ounces per 100 pounds of seed as a planter box treatment.

WHEAT - Use at the rate of 3 ounces per bushel or 4½ ounces per 100 pounds of seed as a drill box treatment. This treatment is not recommended for control of Bunt (Smut).

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label information.

NET WEIGHT

Manufactured By

10 POUNDS

HOPKINS AGRICULTURAL CHEMICAL CO.

MADISON, WISCONSIN 53701

HACCO .5.9.74

