

Crop	Insect Controlled	Rate of Application	Pre-Harvest Interval	Application Timing and Miscellaneous
	Lesser Cornstalk Borers	20-30 lbs. / A	21 days	Apply in band over fruiting zone just before pegging. Cover band with 1-2 inches of soil. Livestock (1) may be fed peanut hulls, (2) may be fed peanut forage 7 days following treatment, and (3) may be fed peanut hay 21 days following treatment.
Sorghum	Cutworms (surface and subterranean)	20-30 lbs. / A	7 days (grain harvest). Crop may be fed to livestock immediately.	Broadcast just before planting. Work into soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Lesser Cornstalk Borers	10-20 lbs. / A	7 days (grain harvest). Crop may be fed to livestock immediately.	Apply in 10-inch band over row at planting or as corn emerges. Cover band with 1-2 inches of soil.
Soybeans	Cutworms (surface and subterranean)	20-30 lbs. / A	—	Apply just before planting. Work into soil immediately, 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Lesser Cornstalk Borers	10-20 lbs. / A	—	Apply in 6-inch band over row just before planting or when soybeans are 1-2 inches high. Cover with 1-2 inches of soil.
Sugar Cane	Sugarcane Borers	15-25 lbs. / A	28 days	Broadcast after joints begin to form and when at least 5% of plants show borers feeding in or under leaf sheath but not bored into stalk. Repeat if necessary. Do not make more than 4 applications per growing season. Bagasse and forage may be fed to livestock 28 days after last application.
Sugar Cane	Wire Worms	60 lbs. A	—	Apply in 12-inch band centered over furrow directly to seed pieces at covering time.
Tobacco	Cutworms (surface or subterranean)	20-30 lbs. A	—	Broadcast before transplanting. Work immediately into soil 2-4 inches.
	Seed Corn Maggots	15 lbs. A	—	Broadcast 3-4 weeks before transplanting and work into soil 2 inches immediately.
	Wire Worms	20-30 lbs. A	—	Broadcast 3-4 weeks before transplanting and work into soil 6-9 inches immediately.
Turf Insects				
Grass Turf	Lawn Chinch Bugs	1-2 lbs. 1000 Sq. Ft.	—	For granular formulations, brush lawn to work granules down to soil surface and water lawn thoroughly after application. Also repeat application in 7-10 days if necessary. Keep children and pets off treated areas until the following afternoon.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**HOPKINS AGRICULTURAL CHEMICAL CO.
MADISON, WISCONSIN 53701**

Hopkins

5% DIAZINON^{®*} GRANULAR INSECTICIDE

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 5%

INERT INGREDIENTS: 95%

TOTAL 100%

* Diazinon Registered Trademark of Ciba-Geigy
EPA Reg. No. 2393-279-AA

EPA Est. No. 2393-WI-1

CAUTION

KEEP OUT OF REACH OF CHILDREN


Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin.

FISH AND WILDLIFE CAUTION:

This product is toxic to fish, shrimp, crabs, birds, and other wildlife. Keep out of lakes, streams or ponds. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

DO NOT RE-USE THIS CONTAINER - DESTROY WHEN EMPTY

Keep children and pets off treated areas until this material is washed into the soil and grass is dry



Manufactured By

HOPKINS AGRICULTURAL CHEMICAL CO.

MADISON, WISCONSIN 5370

NET WEIGHT

POUNDS



5% DIAZINON[®] GRANULAR INSECTICIDE

Direction for Use

Applications of Hopkins 5% Diazinon[®] Granular Insecticide should be applied as per the following recommendations

Crop	Insect Controlled	Rate of Application	Pre-Harvest Interval	Application Timing and Miscellaneous
Strawberries	Mole Crickets	20 lbs. A	5 days	Apply when insects present. Apply 1-2 days before transplanting and mix into top 1-2 inches of soil immediately.
Tomatoes	Vinegar Flies	10-20 lbs. A	0 days	Broadcast when flies first appear. Repeat in 7-10 day intervals or as required.
Carrots	Carrot Rust Flies	40 lbs. A	0 days	Apply to seed furrow at planting.
Broccoli, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Cucumbers, Endive (escarole), Kale, Lettuce, Lima Beans, Muskmelons, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Sweet Corn, Swiss Chard, Tomatoes, Watermelons, Winter Squash	Cutworms	40-80 lbs. A 40-60 lbs. A on Sweet Corn	0 days 0 days	Apply broadcast just before planting. Work into soil. Immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms
Broccoli, Cabbage, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Tomatoes, Turnips	Mole Crickets	20 lbs. A	0 days	Apply to plant beds or fields 1-2 days before sowing seed or transplanting. Work into soil 1-2 inches immediately.
Onions	Onion Maggots	20 lbs. A	0 days	Apply in furrow with seeds at planting.
Broccoli, Peas, Spinach, Cabbage, Cauliflower	Root Maggots	40-60 lbs. A	0 days	Apply broadcast and immediately work into soil 3-4 inches before seeding, either in seed bed or before transplanting.
Radishes	Root Maggots	30-40 lbs. A	--	Apply broadcast and immediately work into soil 3-4 inches before seeding, either in seed bed or before transplanting.

Crop	Insect Controlled	Product Application	Pre-Harvest Interval	Application Timing and Miscellaneous
Turnips, Cabbage, Cauliflower Broccoli	Root Maggots	20 lbs. / A	—	Apply in seed furrow at planting.
Broccoli, Cabbage, Cantaloupes, Carrots, Cauliflower, Celery, Collards, Cucumbers, Endive, (escarole), Kale, Lettuce, Lima Beans, Muskmelons, Onions, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Corn, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelons, Winter Squash	Wire Worms	40-80 lbs. A	—	Apply broadcast before planting and work into soil 4-8 inches immediately.
Potatoes	Wire Worms	20-40 lbs. A	0 days	Apply at planting placing Diazinon in bands 3-6 inches deep along both sides of row.
Corn (Field and Sweet)	Corn Rootworm Larvae	25-50 oz. per 1000 Linear Feet	Corn may be harvested immediately following last application. Do not feed treated forage to dairy or beef cattle or sheep within 10 days after foliar application exceeding 1 1/4 lbs. active ingredient per acre	Apply in 7 inch band centered over corn row at planting.
Corn (Field and Sweet)	Cut Worms	40-60 lbs. A	Corn may be harvested immediately following last application. Do not feed treated forage to dairy or beef cattle or sheep within 10 days after foliar application.	For surface cutworms apply just before planting or when larvae first appear and work into soil 1-2 inches immediately. Do not apply later than layby. For subterranean cutworms apply just before planting and work into soil 3-6 inches immediately.
Corn (Field and Sweet)	Wire Worms	40-80 lbs. per Acre	Corn may be harvested immediately following last application. Do not feed treated forage to dairy or beef cattle or sheep within 10 days after foliar application.	Apply before planting and work into soil 4-8 inches immediately.
Corn (Field and Sweet)	European Corn Borers	20-40 lbs. A	Corn may be harvested immediately following last application. Do not feed treated forage to dairy or beef cattle or sheep within 10 days after foliar application.	Apply so granules fall into whorls when 75% of plants show first generation borers feeding. Apply second application 7 days later if needed. For second generation borers apply when field count shows 10 or more masses per 100 plants. Application may be made by aircraft.