

DIRECTIONS

The following general directions are supplied by us gratuitously but are in no way guaranteed.

For use on non-crop land such as Right-of-Ways, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas and Parking Lots for control of most annual and perennial grasses and broad-leaved weeds. Control will usually last 8 months or longer. The effectiveness and length of time control will last depend on type of soil, amount of rain and other conditions.

RATE: Under ordinary conditions use at the rate of 200 to 400 pounds per acre. Under conditions where weeds are difficult to control increase the rate to 600 pounds per acre. For spot treatment, use 1/2 to 1 pound per 100 square feet. For uniform control spread the material evenly over the area to be treated.

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated, or sift directly from this shaker top can. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control. Do Not apply when ground is frozen.

FLORIDA CITRUS GROVES

For Use in Drainage Ditches For Control of Most Annual and Perennial Grasses and Broad-Leaved Weeds such as Paragrass, Arrowhead, Guineagrass and Water Primrose.

Rate: Use at the rate of 200 to 400 pounds per acre. (See chart)

Amount to Apply Per 100 Linear Feet of Ditch					
Width of Ditch (feet)	2	4	6	8	10
Pounds Urox 11	1 to 2	2 to 4	3 to 6	4 to 8	5 to 10

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. Treat only bottom not banks.

WHEN TO APPLY: Apply when ditch is not in use but soil is moist. If ditch bottom is not moist, wet it down before treating. If less than 4 inches of rain falls between time of treatment and time of intended use, run water into the ditch and let stand for 3 days. Drain off and waste the water before using ditch.

NOTE: Do Not use in ditches into which tree or other desirable plant roots extend. Do Not contaminate domestic water supplies, farm ponds or water to be used for livestock.

CAUTION!

**HARMFUL IF SWALLOWED
IRRITATING TO SKIN AND EYES**

Do not get on skin, in eyes or on clothing. In case of contact, immediately wash skin with plenty of water: for eyes, wash with water for at least 15 minutes and get medical attention.

NOTE: Do not allow to drift onto desirable plants or cropland. Do not apply near lawns, valuable trees or other desirable plants or to areas where their roots may extend. Equipment used for application should be thoroughly cleaned to prevent corrosion or contamination of other materials used in it later.

Seller makes no warranty of any kind, express or implied, concerning the results obtained by the use of this material. Buyer assumes all risk of use or handling whether or not in accordance with any suggestions, recommendations or directions of seller.



ACCEPTED

Apr 19, 1963

REGISTERED TRADEMARK OF ALLIED CHEMICAL COMPANY
FOR ECONOMIC INSECTICIDES
ED. UNDER NO. 2393-242
TO ATTACHED COMMENTS

UROX

11

WEED KILLER

For General Weed Control
on Non-Crop Lands

CODE 9941

ACTIVE INGREDIENT:
Monuron Trichloroacetate [3-(p-chlorophenyl)-1,1-dimethylurea trichloroacetate] 11%
INERT INGREDIENTS: 89%

U. S. PATENT NO. 2,782,112



DIRECTIONS

The following general directions are supplied by us gratuitously but are in no way guaranteed.

For use on non-crop land such as Right-of-Ways, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas and Parking Lots for control of most annual and perennial grasses and broad-leaved weeds. Control will usually last 8 months or longer. The effectiveness and length of time control will last depend on type of soil, amount of rain and other conditions.

RATE: Under ordinary conditions use at the rate of 200 to 400 pounds per acre. Under conditions where weeds are difficult to control increase the rate to 600 pounds per acre. For spot treatment, use 1/2 to 1 pound per 100 square feet. For uniform control spread the material evenly over the area to be treated.

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated, or sift directly from this shaker top can. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control. Do Not apply when ground is frozen.

FLORIDA CITRUS GROVES

For Use in Drainage Ditches For Control of Most Annual and Perennial Grasses and Broad-Leaved Weeds such as Paragrass, Arrowhead, Guineagrass and Water Primrose.

Rate: Use at the rate of 200 to 400 pounds per acre. (See chart)

Amount to Apply Per 100 Linear Feet of Ditch					
Width of Ditch (feet)	2	4	6	8	10
Pounds Urox 11	1 to 2	2 to 4	3 to 6	4 to 8	5 to 10

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. Treat only bottom not banks.

WHEN TO APPLY: Apply when ditch is not in use but soil is moist. If ditch bottom is not moist, wet it down before treating. If less than 4 inches of rain falls between time of treatment and time of intended use, run water into the ditch and let stand for 3 days. Drain off and waste the water before using ditch.

NOTE: Do Not use in ditches into which tree or other desirable plant roots extend. Do Not contaminate domestic water supplies, farm ponds or water to be used for livestock.

CAUTION!

**HARMFUL IF SWALLOWED
IRRITATING TO SKIN AND EYES**

Do not get on skin, in eyes or on clothing. In case of contact, immediately wash skin with plenty of water: for eyes, wash with water for at least 15 minutes and get medical attention.

NOTE: Do not allow to drift onto desirable plants or cropland. Do not apply near lawns, valuable trees or other desirable plants or to areas where their roots may extend. Equipment used for application should be thoroughly cleaned to prevent corrosion or contamination of other materials used in it later.

Seller makes no warranty of any kind, express or implied, concerning the results obtained by the use of this material. Buyer assumes all risk of use or handling whether or not in accordance with any suggestions, recommendations or directions of seller.



ACCEPTED
 Apr 19, 1963
 REGISTERED FEDERAL INSECT
 FUNGICIDE AND RODENT BAIT
 FOR ECONOMIC PEST CONTROL
 ED UNDER NO. 2393-242
 TO ATTACHED DOCUMENTS

ROX[®] 11 WEED KILLER

For General Weed Control
on Non-Crop Lands

CODE 9941

INGREDIENT:
 Monuron Trichloroacetate [3-(p-chlorophenyl)-1,
 1-dimethylurea trichloroacetate] 11%
INGREDIENTS: 89%

U. S. PATENT NO. 2,782,112

DIRECTIONS

The following general directions are supplied by us gratuitously but are in no way guaranteed.

For use on non-crop land such as Right-of-Ways, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas and Parking Lots for control of most annual and perennial grasses and broad-leaved weeds. Control will usually last 8 months or longer. The effectiveness and length of time control will last depend on type of soil, amount of rain and other conditions.

RATE: Under ordinary conditions use at the rate of 200 to 400 pounds per acre. Under conditions where weeds are difficult to control increase the rate to 600 pounds per acre. For spot treatment, use 1/2 to 1 pound per 100 square feet. For uniform control spread the material evenly over the area to be treated.
HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated, or sift directly from this shaker top can. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control. Do Not apply when ground is frozen.

FLORIDA CITRUS GROVES

For Use in Drainage Ditches For Control of Most Annual and Perennial Grasses and Broad-Leaved Weeds such as Paragrass, Arrowhead, Guineagrass and Water Primrose.

Rate: Use at the rate of 200 to 400 pounds per acre. (See chart)

Amount to Apply Per 100 Linear Feet of Ditch					
Width of Ditch (feet)	2	4	6	8	10
Pounds Urox 11	1 to 2	2 to 4	3 to 6	4 to 8	5 to 10

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. Treat only bottom not banks.

WHEN TO APPLY: Apply when ditch is not in use but soil is moist. If ditch bottom is not moist, wet it down before treating. If less than 4 inches of rain falls between time of treatment and time of intended use, run water into the ditch and let stand for 3 days. Drain off and waste the water before using ditch.

NOTE: Do Not use in ditches into which tree or other desirable plant roots extend. Do Not contaminate domestic water supplies, farm ponds or water to be used for livestock.

CAUTION!

**HARMFUL IF SWALLOWED
 IRRITATING TO SKIN AND EYES**

Do not get on skin, in eyes or on clothing. In case of contact, immediately wash skin with plenty of water: for eyes, wash with water for at least 15 minutes and get medical attention.

NOTE: Do not allow to drift onto desirable plants or cropland. Do not apply near lawns, valuable trees or other desirable plants or to areas where their roots may extend. Equipment used for application should be thoroughly cleaned to prevent corrosion or contamination of other materials used in it later.

Seller makes no warranty of any kind, express or implied, concerning the results obtained by the use of this material. Buyer assumes all risk of use or handling whether or not in accordance with any suggestions, recommendations or directions of seller.



SPECIMEN LABEL

UROX[®] 11 WEED KILLER

For General Weed Control
on Non-Crop Lands

CODE 9941

ACTIVE INGREDIENT:
 Monuron Trichloroacetate [3-(p-chlorophenyl)-1,
 1-dimethylurea trichloroacetate] 11%
INERT INGREDIENTS: 89%

U. S. PATENT NO. 2,782,112

