

BEST DOCUMENT AVAILABLE

U

Hopkins

ACCEPTED
MAY 9 1971

UROX[®] 11

WEED KILLER

For General Weed Control on Non-Crop Lands

ACTIVE INGREDIENTS:	
Mecuron (3,7-dimethyl-2,6-dimethyl-4-pyridinylmethyl dimethylcarbamate)	11
1,1-dimethyl-2-methylpropylamine hydrochloride	6.9
INERT INGREDIENTS	82.1
Total	100

EPA Reg. No. 100-100000-0100
EPA Reg. No. 100-100000-0100
EPA Reg. No. 100-100000-0100

CAUTION

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
IRRITATING TO SKIN AND EYES**

Do not get on skin, in eyes or on clothing. In case of contact, immediately wash skin with plenty of water; for eyes, wash with water for at least 15 minutes and get medical attention.

NOTE: Do not use on areas where roots of desirable plants extend.

Do not apply to areas where roots of desirable plants or to areas where their roots may extend. Equip sprayer with filter to prevent corrosion or contamination of other materials.

DIRECTIONS

NON-CROP LAND

For use on non-crop land such as Rights-of-way, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas, Parking Lots and Under Asphalt for control of annual and perennial grasses and broadleaved weeds. Control will last for 6 to 8 weeks or longer. The effective speed and length of time control will last depend on type of soil and other conditions.

RATE: For ordinary conditions use at the rate of 200 to 400 pounds per acre. Under conditions where weeds are difficult to control, increase the rate to 600 pounds per acre. For spot treatment use 1 to 1.5 pounds per 100 square feet. Broadcast and spread the material evenly over the area to be treated.

HOW TO APPLY: Use a mechanical sprayer or other device which will spread the required amount evenly over the area to be treated. May be applied when top growth is 1 to 2 inches tall, but better control will be obtained when weeds are 3 to 4 inches tall. Excess growth is rapid. Consider before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or irrigation after application will cause the material to move to the root zone and thus speed up control. Do not apply when ground is frozen.

FLORIDA CITRUS GROVES

For Use in Drainage Ditches For Control of Most Annual and Perennial Grasses and Broadleaved Weeds such as Paragrass, Arrowhead, Guinea Grass and Water Primrose

RATE: Use at the rate of 200 to 400 pounds per acre. (See chart)

Width of Ditch (feet)	Amount to Apply per 100 Linear Feet of Ditch				
	2	4	6	8	10
1 to 2	1 to 2	2 to 4	3 to 6	4 to 8	5 to 10

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. Treat only bottom, not banks.

WHEN TO APPLY: Apply when ditch is not frozen but soil is moist. If ditch bottom is not moist, wet it down before treating. Allow 4 to 6 feet of topsoil between time of treatment and time of intended use. Run water into the ditch and let stand for 3 days. Drain off and waste the water before using ditch.

NOTE: Do Not use on drainage ditches to which tree or other desirable plant roots extend. Do Not contaminate domestic water supplies, farm ponds, or water to be used for livestock.

NOTICE OF WARRANTY

Read directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals, and specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.



NET WEIGHT

50 POUNDS