

FERBAM

DIRECTIONS

The following general directions are supplied by us gratuitously and are believed to be reliable, but are in no way guaranteed.

MIXING SPRAYS: Wash Ferbam into spray tank through a deep screen, or make a thick paste with a little water, then dilute it and add to spray tank while filling.

SPREADER-STICKER: The addition of Orchard

Brand Filmfast®, Genifilm® "L", or Plyac® will improve coverage on foliage which is difficult to wet.

Dosages below are based on the amount of Ferbam to be used in 100 gallons of water.

FRUIT

APPLES: For Apple Scab, use 1 to 1½ pounds. Apply in pre-bloom, calyx and early cover sprays. Reduce to ¾ pound in later cover sprays. When used in combination with sulfur, Ferbam is recommended at ½ pound, plus 3 pounds of micron-fine wet-table sulfur (Orchard Brand Micro-Dritomic® Sulfur). Do Not use sulfur if russetting has been a problem. Do Not use Ferbam on tender varieties such as Golden Delicious as it may contribute to fruit russetting.

For Cedar Apple Rust Diseases, keep foliage and fruit covered during infection periods. Use as above, plus a special bloom spray of ½ pound.

For Black Rot (Frog-eye) (Leaf and Fruit Spot), Apple Blotch, Bitter Rot, Sooty Blotch and Fly Speck, use 1 to 1½ pounds.

CHERRIES: For Leaf Spot, use 1½ pounds. Apply 4 sprays, at petal-fall, 2 and 4 weeks after petal-fall and immediately after harvest. For Brown Rot, use 1½ pounds. Apply at popcorn, full-bloom and petal-fall, and starting 2 to 3 weeks before harvest apply one or more sprays at 7 to 10 day intervals.

PEARS: For Scab, use 1½ pounds. Apply in pink, calyx, first cover spray and second cover spray. For Fabraea (Leaf Blight and Fruit Spot) and Sooty Blotch, use 1 pound in the summer sprays.

PEACHES: For Peach Leaf Curl and Shot Hole (fungus), use 1½ to 3 pounds. Apply at dormancy and again in early bloom. For Brown Rot Blight, use 1½ to 2 pounds. Apply at pink bud and 25% to 75% of full bloom. For Fusilcoccum Canker (constriction) in New Jersey and Massachusetts, use 1½ pounds. Apply before bloom (starting in early March) and after harvest as long as weather permits. Space sprays 7 to 10 days apart.

GRAPES: For Black Rot, use 2 pounds. Apply just before bloom, immediately after bloom and 2 weeks after bloom. Do Not use more than 6 lbs. per acre per application.

CRANBERRIES: For Fruit Rots, use 3 pounds. Apply at mid-bloom and repeat at 2 week intervals, for a total of 3 applications. Do Not use more than 9 pounds per acre per application.

NOTE: Do Not use on apples, pears, or grapes later than 7 days before harvest; on peaches Not later than 21 days before harvest; on cranberries Not later than 28 days after mid-bloom.

OTHER CROPS

CABBAGE: For Downy Mildew in plant beds, use 1½ pounds. Make 2 to 3 applications a week starting when plants emerge. Do Not use more than 3 pounds per acre per application.

TOBACCO: For Blue Mold in plant beds, use at the rate of 3 to 4 pounds per 100 gallons of water. Use the 4-pound rate if equipment and timing are not perfect. Apply 3 gallons per 100 square yards when plants are small. Increase to 6 gallons as plants are ready for transplanting. Stop applications when leaves are the size of a dime or when Blue Mold first appears in your area. Apply twice a week until time for transplanting.

ORNAMENTALS: For Black Spot and Rust of Roses; Septoria Leaf Spot and Rust of Chrysanthemums; Snapdragon Rust and

Leaf Spot; Bacterial Leaf Spot of Geraniums; Alternaria Blight and Rust of Carnations; Leaf Spot and Botrytis Rot of Cyclamen; Botrytis Blight (tulip fire) of Tulips; Juniper Rust, and Aster, Begonia, Gardenia, Hydrangea, Ivy, and Laurel Leaf Spots—Use 1 to 1½ pounds. Apply at 7 to 10 day intervals as needed. For Petal Blight of Azaleas and Camellias—use 1 to 1½ pounds. Apply while buds are opening and repeat at 3 to 5 day intervals as needed during infection periods.

SMALL QUANTITIES: Use at the rate of 4, 6 or 8 teaspoonfuls in 1 gallon of water, or equivalent to 1, 1½ or 2 pounds in 100 gallons of water.

Seller makes no warranty of any kind, express or implied, concerning results obtained by the use of this material. It is assumed all use or handling whether or not in accordance with any suggestions, recommendations or directions of seller.

CAUTION: MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN.

Avoid breathing dust or spray mist.

Avoid contact with eyes, skin or clothing.

In case of contact, flush with plenty of water.

For eyes, get medical attention.

Keep away from fire and sparks.