

Hopkins 25% Lindane Wettable Powder

EPA Reg. No. 2393-305

Paper & Plastic Bag

Center Panel

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision
and only for those uses covered by the Certified Applicator's certification.

Hopkins

25% LINDANE

WETTABLE POWDER

For Use on Fruit and Vegetables


For Use on Livestock

ACTIVE INGREDIENT LINDANE (Gamma Isomer of Benzene Hexachloride) 25%
INERT INGREDIENTS 75%


KEEP OUT OF REACH OF CHILDREN

WARNING

EPA Reg. No. 2393-305
EPA Est. No.



For Additional Warning Information See Other Panel



NET WEIGHT

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed.

May be absorbed through skin.

Do not breathe dust, vapor or spray mist.

Do not get on skin or in eyes.

Do not contaminate feed and foodstuffs and water supplies.

Wash thoroughly after handling and before eating or drinking.

Applicators must wear the following protective clothing during the application process: a lightweight protective suit or coveralls; water-resistant hat; unlined, waterproof gloves; and unlined, lightweight boots.

Do not allow children to handle or apply this product. Children and pets should not be allowed in treated areas until sprayed surfaces are dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not contaminate any body of water. Do not use where runoff is a frequent problem. Birds feeding in treated areas may be killed.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Do not apply when weather conditions favor drift from the treated area.

NOTICE OF WARRANTY

Follow directions carefully. Results may vary due to application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other factors. Factors in the use of this product are the responsibility of the seller. Buyer assumes all risks of injury, damage or handling of this material not mentioned herein, and with directions given hereon.

DIRECTIONS FOR USE

It is a violation of Federal law
to use this product in a manner
inconsistent with its labeling.

All dosages shown below are based
upon the amount of 25% Lindane Wettable
Powder required per 100 gallons of
water.

Avoid use of this material where soil may become contaminated as this
may cause crop injury. Do not use more
than the following amounts of this material per acre per application:
to foliage, on strawberries 4 pounds, on celery and on leeks 3 pounds, on
peppers 2 1/2 pounds, and on all other vegetable crops listed 3 pounds.
Do not use more than 4 pounds for soil treatment for cutworms and
caterpillars. **DO NOT APPLY BY AIRCRAFT**

VEGETABLES AND FRUITS

APPLES For Aphids, Leafhoppers and Grasshoppers use 1 pound when
insects are present. Do not apply on Apples within 60 days of harvest.
Do not apply after first frost as off flavor may result. Do not use
more than 14 pounds per acre.

CUCUMBERS For Aphids, Thrips, Sawflies, Leafhoppers, Caterpillars
and other insects use 1 pound when insects are present.
Do not use after Cucumbers are full grown. Do not apply after Cucumbers start to bunch.
Use 100 to 150 gallons per acre.

EDIBLE CROPS (Cabbage and Broccoli) For Aphids, Flea Beetles, Caterpillars
and Thrips use 1 pound when insects are present. Use
100 to 150 gallons per acre. For Cabbage Mosaic use 4 pounds per
acre in enough water to give 1/2 inch coverage. Apply to soil around base
of plants. Use only before heads start to form.

CUCUMBERS (Cucumber and Squash) For Cucumber Beetles and Thrips
use 1 pound. For Leafhoppers, Flea Beetles, Caterpillars, Flea Bugs,
and Aphids use 1 pound. Use only before Cucumbers start to bunch and when in
soil are present. Use 100 to 150 gallons per acre.

CHICKENS For Thrips use 1 pound when Thrips are present. Use 100 to
150 gallons per acre. For Leafhoppers use 1 pound when insects are present. Use 100 to
150 gallons per acre. For Cabbage Mosaic use 4 pounds per acre in enough
water to give 1/2 inch coverage. Apply to soil around base of plants.
Do not use after Cucumbers start to bunch and when in soil are present. Use 100 to 150 gallons per acre.

PEPPERS For Aphids, Leafhoppers and Armyworms use 1 pound. Do not
apply after Cucumbers start to bunch. Use 100 to 150 gallons per
acre.

SQUASH For Aphids, Leafhoppers and Armyworms use 1 pound. Use only in seedling
stage. Use 100 to 150 gallons per acre.

SWEETPEPPERS For Weevils and Flea Beetles use 1 pound in Spring
when insects appear. Do not apply after Cucumbers start to bunch. Use 100
to 150 gallons per acre.

PEACHES (for fresh fruit market only) For Aphids and Thrips use 1/2
pound when insects are present. For Leafhoppers use 1 pound when
insects are present. Do not apply after Cucumbers start to bunch.
Use 100 to 150 gallons per acre.

COMPATIBILITY Do not use with general insecticides. For details of use of any
other material.

LIVESTOCK

Do not use on animals for more than 3 months or daily animals or
poultry.

Do not spray animals within 30 days of slaughter or dip within 30 days
of slaughter.

Avoid contamination of feed, trough, drinking water and feed troughs.
If drinking or feed troughs are contaminated, wash out IMMEDIATELY
with plenty of water before spraying. Do not use in cold weather
when there is danger of animals contracting respiratory diseases, such
as colds or pneumonia.

ON BEEF, CATTLE, SHEEP, AND GOATS FOR MANGE MITE (SCABIES)
Use 1 pound per 100 lbs. of water. For horses and cattle, use 2
pounds of spray. Do not use on animals with open wounds. Water
animals should be clean for treatment. For mite control use control
agents two weeks apart using sufficient water to part the
animal's hair and wet its skin. Do not treat horses within 2 weeks of
turning out for at least 3 weeks thereafter.

Dip solutions must be disposed of in accordance with the
Resource Conservation and Recovery Act (RCRA). If the applicator
generates more than 1000 kg (2200 lbs.) of used dip solution
per month or more than 1000 kg (2200 lbs.) used dip solution in
combination with other hazardous waste, the material must be
treated as a hazardous waste subject to subpart C of RCRA. Any
user who wishes to treat, store or dispose of hazardous waste,
must obtain a permit to serve as a hazardous waste facility
permitted to RCRA.

DO NOT USE ON WASTE AREA VEGETATION. Use 2 1/2 pounds per
100 gallons when 1/2 inch of growth is present. Do not be used
depends on size of vegetation. Do not permit livestock to graze on
treated areas.

STORAGE AND DISPOSAL 1-12 pt.

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container tightly closed. Store in original container. Protect from direct contact with water or excessive moisture. Once opened, do not store bag for prolonged period.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.