

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMAN & DOMESTIC ANIMALS**

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In both cases, intravenous and oral administrations of Vitamin K<sub>1</sub> combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

**ENVIRONMENTAL HAZARD**

Keep Out of Lakes, Streams or Ponds.


EPA Reg No 2393-363  
EPA Est No 2393-IL-3<sup>A</sup>, 2393-WI-1<sup>A</sup>,  
523-IL-1<sup>A</sup>, 2393-WI-2<sup>A</sup>, 33509-MO-1<sup>A</sup>  
Superscript refers to first letter of lot number  
CODE 2393-363 1 2 3 4 5 6 7 8 9 (8/80)

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings causes fatal hemorrhage in rats and mice. Bacteria which produce Vitamin K<sub>1</sub>, an antidote for Warfarin, are inhibited through an antibacterial agent contained in this product.

**NOTICE OF WARRANTY**

SELLER AND MANUFACTURER MAKE NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL.





**PROLIN<sup>®</sup>**  
*Pelleted Rat Bait*

**ACTIVE INGREDIENTS:**  
Warfarin [3-(alpha-Acetyloxybenzyl)-4-hydroxycoumarin] ..... 0.025%  
N<sup>1</sup>-(2-Quinoxaliny) sulfanilamide (Sulfiquinoxaline) ..... 0.025%  
**INERT INGREDIENTS:** ..... 99.950%  
**TOTAL** ..... **100.000%**

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS**

® PROLINIS MANUFACTURED UNDER LICENSE FROM THE WISCONSIN ALUMNI RESEARCH FOUNDATION.

**NET WEIGHT 1 LB. (0.454 Kg.)**

2393 363  
**Hopkins**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:** For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper proof bait boxes or locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** After removing all food, water, and harborage, as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners, and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or snow.

**APPLICATION DIRECTIONS**

**RATS:** Apply 4 to 6 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply 1/4 to 1/2 ounce (2 level tablespoons) of bait at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. Repeat treatment when infestation recurs. Where continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**DISPOSAL**

DO NOT REUSE EMPTY CONTAINER.  
WRAP CONTAINER AND PUT IN TRASH COLLECTION.