

1/11/4

2393 363

1/2

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN — If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

EPA Reg. No. 2393-363

EPA Est. No. 2393-IL-3[®], 2393-WI-1[®],
523-IL-1[®], 2393-WI-2[®], 33609-MO-1[®]
Superscript refers to first letter of lot number.
CODE 2393-363 1 2 3 4 5 6 7 8 9 (6-82)

PRODUCT DESCRIPTION

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rats and mice.

ENVIRONMENTAL HAZARD

Keep Out of Lakes, Streams or Ponds.

NOTICE OF WARRANTY

SELLER AND MANUFACTURER MAKE NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL.

NET WEIGHT

4 OZ. (114 grams)



Hopkins
MANUFACTURED BY
Hopkins Agricultural Chemical Co.
Waukesha, Wisconsin, U.S.A.

PROLIN IS MANUFACTURED UNDER
LICENSE FROM THE WISCONSIN ALUMNI
RESEARCH FOUNDATION.

See Back Panel for Additional
Precautions

**KEEP OUT OF REACH OF
CHILDREN
CAUTION**

ACTIVE INGREDIENT:
Warfarin (3-(alpha-Acetyloxybenzyl)-
4-hydroxycoumarin) 0.025%
INERT INGREDIENTS: 99.975%
TOTAL 100.000%
Contains 0.025% Sulfadiazine



Hopkins
PROLIN[®]
Pelleted Rat Bait

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS. For control of Norway rats, Roof rats and House mice in and around homes, buildings, and similar man-made structures, and in Transportation Vehicles.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS — Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

READY MIXED, EASY TO USE — USE INDOORS AND OUTDOORS

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

**DO NOT REUSE EMPTY CONTAINER.
WRAP CONTAINER AND PUT IN TRASH COLLECTION.**

BEST AVAILABLE COPY

REGISTERED

ACCEPTED
JAN 24 1983
Office of Federal Registration of
Fungicides and Rodenticides Act
as amended, for the pesticide
registered under
EPA Reg. No. 2393-363

BEST AVAILABLE COPY