

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife.

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARD

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply in areas where animals have received direct application of DDVP within 8 hours.

FOR DAIRY CATTLE, BEEF CATTLE AND HORSES: To protect livestock from face flies, houseflies, stable flies, horse flies, mosquitoes and gnats, spray the animal daily using one to two ounces per head. Cover the animal with spray with particular attention to the back, flanks, legs and shoulders. Do not use in excess of 2 ounces per adult animal. Apply proportionately less to smaller animals. Do not soak the skin or wet the hides. Care should be taken that the spray does not come in direct contact with the lactating dairy cow's teats unless they are washed with an approved cleansing solution and dried before milking. Brahman and Brahman cross cattle should not be treated as they may show hypersensitivity to organic phosphates.

FOR BARN, MILK ROOMS AND STABLES: When used as a space spray, direct power sprayer toward upper portion of the enclosure until area is filled with a fine mist. If a coarse spray is used, apply wherever flies congregate or alight, until surfaces are moistened. Avoid contact with food or milk utensils. Close doors and windows before treatment when possible. This treatment will kill resting adult flies. Do not remain in treated areas and ventilate the area after treatment is complete.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not use or store near heat or open flame. Keep product in original container. Keep container tightly closed. High temperatures may shorten shelflife of product.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



Cabbage (7) Cauliflower (5) Broccoli (5) Radishes (10)	Aphids, Flea Beetles, Imported Cabbage Worm, Dipterous Leafminers
Carrot (10) Lettuce (10)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminer, Mites
Cantaloupes (3) Cucumbers (7) Squash (3)	Aphids, Thrips, Mites
Tomatoes (1) Eggplants (7)	Aphids, Fall Armyworm (tomato), Flea Beetles, Leafhoppers, Southern Armyworm, Vinegar Flies (tomato)
Onions (10)	Onion Thrips
Sweet Corn (1)	Aphids, Flea Beetles, Grasshoppers, Mites
Vegetables: Directions for Use	2 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden. Thoroughly spray all plant parts. Generally, if infestations warrant, application at 7-10 day intervals may be required. Do not feed treated vines to livestock.
VEGETABLES	
Preplant Soil Application	
Carrots, Corn, Lettuce	Cutworms (also radishes and cabbage) 12 fl. oz. per 1,000 sq. ft. Work into 2-3 inches of soil for surface cutworms; 3-6 inches for subterranean cutworms. Wireworms (except red beet) 6 fl. oz. per 500 sq. ft. of soil working into soil 4-8 inches. Root Maggot (cabbage only) 9 fl. oz. per 1,000 sq. ft. Work into soil 4-8 inches. Apply broadcast just before planting. Mix with enough water to achieve thorough coverage and work into soil immediately.
Peas, Red Beans, Soy Beans	
FRUITS & NUTS	
Apples (14) Pears (14) Peaches (7) Nectarines (10)	Number following crops refers to preharvest interval (in days) following spray application. Aphids, Leafing Moths, Fruit Tree Leaf Miner, Maggot (apples), Mites (European Red, Two Spotted Spider), San Jose Scale (Crawler) (Apples and Pear only) 2 fl. oz. in 3 gals. of water. Apply as full cover spray.
Strawberries (5) Elderberries (7) Blackberries (7) Raspberries (7)	Aphids, Two spotted Mites 2 fl. oz. in 3 gals. water. Apply as full cover spray.
Grapefruit, Lemons (7) Oranges	Citrus Thrips, Scale Crawlers (Citrus Snow Scale, Cottony cushion) 2 fl. oz. in 3 gals. of water. Apply as a full cover spray. Start spraying after blooming. Applications should be made according to infestation of when aphids, mites and crawler strains of scale insects are abundant. If infestations warrant, repeat applications at 7-10 day intervals. 2 1/2 fl. oz. in 3 gals. of water. Apply as a full cover spray. Apply after blooming and again in the summer. Thorough spray coverage of tree trunk limbs and twigs is essential.
Almonds, Filberts Pecans, Walnuts	Aphids, Nut Casebearer (pecans), Scale Crawlers, Spider Mites 1 1/2 fl. oz. in 3 gals. of water. Apply as a full cover spray. Do not apply after hard cover is split.
ROSES AND FLOWERS	
Azalea, Camellia Chrysanthemum, Cassia Gladiolus, Lilac, Phlox, Dentist, Flower	Aphids, Flea Beetle, Dipterous Leafminers, Leafhoppers, Mealybugs, Two Spotted Spider Mite, Thrips, Whiteflies 2 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Wet the plants to the dripping point. Try to hit the undersides of leaves and penetrate dense foliage.
TREES & SHRUBS	
Arborvitae Birch, Boxwood Holly, Juniper, Laurel Maple, Mimosa, Oak Pine, Poplar, Spruce, Yew	Aphids, Bagworms, Leafhoppers, Mealybugs, Scale Crawlers (Cottony cushion, Euonymus, San Jose Scale), Webworms (Juniper, Mimosa, Yew) 2 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage. 1 1/2 fl. oz. for Mimosa Webworms. Apply a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals. Tree Caterpillars, Webworms (Lettucebaster, Fall) 2 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife.
Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARD

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not apply in areas where animals have received direct application of DDVP within 8 hours.

To protect livestock from face flies, houseflies, stable flies, horse flies, mosquitoes and gnats, spray the animal daily using one to two ounces per head. Cover the animal with spray with particular attention to the back, flanks, legs and shoulders. Do not use in excess of 2 ounces per adult animal. Apply proportionately less to smaller animals. Do not soak the skin or wet the hides. Care should be taken that the spray does not come in direct contact with the lactating dairy cow's teats unless they are washed with an approved cleansing solution and dried before milking. Brahman and Brahman cross cattle should not be treated as they may show hypersensitivity to organic phosphates.

When used as a space spray, direct power sprayer toward upper portion of the enclosure until area is filled with a fine mist. If a coarse spray is used, apply wherever flies congregate or alight, until surfaces are moistened. Avoid contact with food or milk utensils. Close doors and windows before treatment when possible. This treatment will kill resting adult flies. Do not remain in treated areas and ventilate the area after treatment is complete.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not use or store near heat or open flame. Keep product in original container. Keep container tightly closed. High temperatures may shorten shelflife of product.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.



CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



249 2
5



Fly-Die



STATEMENT OF PRACTICAL TREATMENT

Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.

Remove victim to fresh air. Apply respiration if indicated.

Flush eyes with plenty of water. Get medical attention immediately.

Empty stomach if material was taken internally. Gastric lavage may be indicated. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Avoid breathing spray mist.

Wash thoroughly with soap and water after handling and before eating or smoking.

Do not contaminate water or foodstuffs.

Do not apply in areas where animals have received direct application of DDVP within 8 hours.

Do not contaminate milk or milk equipment.

Continued on back panel.