



### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED**— Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.

**IF INHALED**— Remove victim to fresh air. Apply respiration if indicated.

**IF IN EYES**— Flush eyes with plenty of water. Get medical attention immediately.

**NOTE TO PHYSICIAN:** Empty stomach if material was taken internally. Gastric lavage may be indicated. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Avoid breathing spray mist.

Wash thoroughly with soap and water after handling and before eating or smoking.

Do not contaminate water or foodstuffs.

Do not apply in areas where animals have received direct application of DDVP within 8 hours.

Do not contaminate milk or milk equipment.

Continued on back panel.

USE

28 1/2 x 7

## ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife.

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

## PHYSICAL OR CHEMICAL HAZARD

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply in areas where animals have received direct application of DDVP within 8 hours.

**FOR DAIRY CATTLE, BEEF CATTLE AND HORSES:** To protect livestock from face flies, houseflies, stable flies, horse flies, mosquitoes and gnats, spray the animal daily using one to two ounces per head. Cover the animal with spray with particular attention to the back, flanks, legs and shoulders. Do not use in excess of 2 ounces per adult animal. Apply proportionately less to smaller animals. Do not soak the skin or wet the hides. Care should be taken that the spray does not come in direct contact with the lactating dairy cow's teats unless they are washed with an approved cleansing solution and dried before milking. Brahman and Brahman cross cattle should not be treated as they may show hypersensitivity to organic phosphates.

**FOR BARNS, MILK ROOMS AND STABLES:** When used as a space spray, direct power sprayer toward upper portion of the enclosure until area is filled with a fine mist. If a coarse spray is used, apply wherever flies congregate or alight, until surfaces are moistened. Avoid contact with food or milk utensils. Close doors and windows before treatment when possible. This treatment will kill resting adult flies. Do not remain in treated areas and ventilate the area after treatment is complete.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Do not use or store near heat or open flame. Keep product in original container. Keep container tightly closed. High temperatures may shorten shelflife of product.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL: Metal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.