

PM-19

Reg # 2393 - 375

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

SSISL76

300

17

194

MAR 21 1997

Glenda Haage
Platte Chemical Company
P.O. Box 667
Greeley, CO 80632-0667

Subject: Hopkins Poultry and Garden Dust
EPA Registration No. 2393-375
Amendment dated December 2, 1996
Response to Agency letter dated 10/1/96

Dear Ms. Haage:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the following changes to your label:
 - a) Delete the following sentences from the PRECAUTIONARY STATEMENTS: "Wash thoroughly after handling. Take shower after work. Change contaminated clothing daily. Wear regular long-sleeved clothing." The correct language now appears in the USER SAFETY REQUIREMENTS box.
 - b) In the Directions for Use on CORN, change "...repeat at 2 to 3 day intervals..." to "...repeat at 7 day intervals..."
2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these

	Conditions	CONCURRENCES					
SYMBOL							
SURNAME							
DATE							

Prohounds A:1293/395.1 3/19/97

274

-2-

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

/s/

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

Hopkins®

394

POULTRY AND GARDEN DUST


96 DEC 11 A 8:46

RECD LAM/OP/P/DPD1

An insecticidal dust for use against certain insects attacking Poultry and Vegetables.

ACCEPTED with COMMENTS in EPA Letter Dated

MAR 21 1997



Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2393-375

ACTIVE INGREDIENT:		
Carbaryl (1-Naphthyl methylcarbamate)	5.00%
INERT INGREDIENTS:	95.00%
	TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 2393-375

EPA EST. NO. 34704-NB-1 (Lot No. Begins 08)

EPA EST. NO. 44616-MO-1 (Other Lot No.)

NET WEIGHT 2 LBS. (907.2 G)

5873

EXP 1/96

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Take shower after work. Change contaminated clothing daily. Wear regular long-sleeved clothing. Avoid contamination of feed and food-stuffs. Do not feed treated vines or trash to meat or dairy animals.

NOTE FOR PHYSICIANS: Carbaryl is a moderate reversible, Cholinesterase inhibitor. Atropine is antidotal.

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. Do not allow children or pets to contact treated areas until dust has settled.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer, or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container only, preferably in a locked area. Keep container tightly closed when not in use. Store in a cool, dry place. High temperatures may shorten shelf life of product. Do not use food or drink containers for mixing or storage. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse empty container. Wrap container and put in trash.

CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS SUCH AS PARTRIDGES AND PHEASANTS, AND PIGEONS:

Litter Treatment: Apply 1 pound per 40 square feet to control Northern Fowl Mite, Chicken Mite, Lice and Bed Bugs. Use a scoop and scatter by hand or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

Hand Application To Birds: Use 1 pound to treat 100 birds by means of a shaker can, squeeze bottle or hand duster to control Northern Fowl Mite, Chicken Mite, Lice and Fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for Chicken Mite and Fleas is a supplement to litter treatment for control of these pests. Do Not apply within 7 days of slaughter.

Poultry Houses: Apply 25 pounds per 1,000 square feet. Apply thoroughly to floors, roosts, and interior surfaces. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter.

VEGETABLES IN HOME GARDENS:

For Mexican Bean Beetles, Leafhoppers, Lygus and Stink Bugs on beans, apply 1 to 2 oz. per 50 row feet. Dust every 7 to 10 days or as needed. May be applied up to harvest at rate directed without

4 7 4

POULTRY AND GARDEN DUST

EPA REG. NO. 2393-375

exceeding residue tolerances.

For **Striped and Spotted Cucumber Beetles, Squash Bugs, Flea Beetles, Pickleworm and Melonworm** on cucumbers, summer squash, apply 1 ounce per 50 row feet. Dust every 7 to 10 days or as needed. May be applied up to day of harvest at rate directed without exceeding residue tolerances.

For **Tomato Fruitworm, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer and Stink Bugs** on tomatoes, apply 1 to 2 oz. per 50 row feet. Dust every 7 to 10 days or as needed. May be applied up to day of harvest at rate directed without exceeding residue tolerances.

For **Colorado Potato Beetle, Leafhoppers, Flea Beetles and Lygus Bugs** on potatoes, apply 1 to 2 oz. per 50 row feet. Dust every 7 to 10 days or as needed. May be applied as foliage application up to harvest at rate directed without exceeding residue tolerances.

For **Corn Earworms, on sweet corn**, apply 1 to 2 oz. per 50 row feet. Begin dusting when first silks appear and repeat at 2 to 3 day intervals until silks are withered. May be applied up to harvest at rate directed without exceeding residue tolerances.

NOTICE OF WARRANTY

SELLER AND MANUFACTURER MAKE NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL.



MANUFACTURED FOR

HACO, inc.

BOX 7190
MADISON, WISCONSIN 53707 U.S.A.