

BEST AVAILABLE COPY

Reg # 2393-408
PM 12
pg 2014

GARDEN DUST

A Combination Fungicide and Insecticide

NEW • IMPROVED • CONTAINS SEVIN®

Active Ingredients:

Maneb (Manganese ethylene bisdithiocarbamate) 3.9%

Carbaryl (1-Naphthyl N-Methylcarbamate) 5.0%

Inert Ingredients (including ZnSO₄) 91.1%

EPA Reg. No. 2393-408

This product is a trademark of the Dow Chemical Company.

A Dual Purpose Product: Highly effective against listed insects and fungi. It is no longer necessary to purchase several different products to give good control in your garden. You can rely on Mohees Garden Dust to safeguard your plants and shrubs effectively.

Economical to Use: Helps protect your investment in your flower and vegetable gardens and ornamental shrubs.

Ready to Use: Needs no mixing or diluting—simply use as it comes from the package.

Highly Effective: Kills listed insects and will control many of the most prominent destructive blights or fungus diseases.

Apply as soon as plants become infested with insects and damage is noticeable. Apply at 5 to 10 day intervals as needed. Dust thoroughly to get good coverage on upper and lower sides of leaves. Use a good dust. Dust adheres better in late evening or early morning when vegetation is moist with dew and when the air is calm.

KEEP OUT OF REACH OF CHILDREN ← 12 pt
CAUTION ← 12 pt

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Avoid contact with eyes and skin. Do not take internally. Skin contact may be harmful.

Avoid skin contact. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD.

DIRECTIONS FOR USE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Keep container tightly closed. Store in original container. High temperatures may shorten shelflife of product.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty container. Discard empty container in trash.

ACCEPTED
with COMMENTS
EPA Letter Dated

APR 29 1985

2393-408

Reg # 2393-408
PM 12
PS 3 of 4Hopkins Garden Dust
EPA Reg. No. 2393-408
Carbaryl L.I.P.
page 2

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s)

BEANS—(Cowpeas, Blackeye Peas)—for Downy Mildew, (Lima, Snap, Green)—for Anthracnose, Downy Mildew, (Cowpeas, Blackeye Peas, Lima, Snap, Green)—for Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Flea Beetles, Japanese Beetles, Leafhoppers, Velvetbean Caterpillars, Western Bean Cutworms, Armyworms, Corn Earworms, Stink Bugs, Tarnished Plant Bugs. Dust plants thoroughly at first sign of disease or infestation. Repeat at 5 to 10 day intervals. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

CARROTS—for Cercospora Leaf Spot, Alternaria, Flea Beetles, Leafhoppers, Harlequin Bugs. Begin when plants are 6 weeks old. Repeat at 7 to 10 day intervals.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER—for Alternaria Leaf Spot, Downy Mildew, Flea Beetles, Harlequin Bugs, Armyworms, Cabbage Worms and Loopers, Corn Earworms. Dust plants thoroughly at first sign of disease or infestation. Repeat at 3 to 7 day intervals. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

SWEET CORN—for Helminthosporium Leaf Blight, Corn Earworms, Corn Rootworm Adults, European Corn Borers, Fall Armyworms, Flea Beetles, Japanese Beetles, Sap Beetles and Leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ear, apply at 3 day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Do not feed treated forage to dairy animals or to animals being fed for slaughter. Applications of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

PEAS—for Downy Mildew, Anthracnose, Cercospora Leaf Spot, Burning Stem Blight, Flea Beetles, Mexican Bean Beetles, Leafhoppers. Begin at late budding when vines begin to run or when disease or infestation first appears. Do not use or watermulsions in Florida. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

CUCUMBERS—Downy Mildew, Anthracnose, Alternaria, Macrostium Leaf Spot, Angular Leaf Spot, Pythium Fruit Rot, Fall Armyworms, Cucumber Beetles, Flea Beetles, Leafhoppers. Begin when plants begin to run or when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

PUMPKINS—for Downy Mildew, Angular Leaf Spot, Flea Beetles, Leafhoppers, Mexican Bean Beetles. Begin when disease or infestation first begins. Repeat at 7 to 10 day intervals.

SQUASH—for Downy Mildew, Anthracnose, Flea Beetles, Leafhoppers, Squash Bugs. Begin when plants start to run or when infestation first appears. Repeat at weekly intervals. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

GRAPES—for Black Rot, Gray Rot, Moths, Japanese Beetles, Flea Beetles, Red Banded Leaf Rollers, Grape Leaf Rollers, Grape Leafhoppers and Grape Leaf Skeletonizers. Begin just before bloom to full bloom. Apply as necessary. Apply just before first brood leaf folder or as one of first releases as needed for Leafhoppers.

COLLARDS—for Downy Mildew, Alternaria Leaf Spot, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Stink Bugs, Tarnished Plant Bugs, Lygus Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

ENDIVE, LETTUCE—Downy Mildew, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Stink Bugs, Tarnished Plant Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

SPINACH—for Downy Mildew, White Rust, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Stink Bugs, Tarnished Plant Bugs, Lygus Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

EGGPLANT—for Anthracnose, Early Blight (Alternaria), Phomopsis Blight or Fruit Rot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when fruit starts to form or when infestation appears. Repeat at 7 to 10 day intervals.

PEPPERS—for Anthracnose, Cercospora Leaf Spot, Ripe Rot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals.

TOMATOES—(Greenhouse and Field) for Cladosporium Leaf Mold, Early and Late Blight, Gray Leaf Spot (Tomphylium), Septoria Leaf Spot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tomato Fruit Worms, Tomato Hornworms, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when fruit forms or infestation appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

ROSES—for Black Spot, Cercospora Leaf Spot, Rust. Begin when first leaves unfold. Repeat at 7 to 10 day intervals.

GRASSHOPPERS on Above Crops—Dust as necessary.

BOX ELDER BUGS—Dust around foundation of house and a few feet of surrounding lawn.

Reg # 2393-408

CM 12

Pg 4 of 4

Hopkins Garden Dust

EPA Reg. No. 2393-408

Carbaryl I.I.P.

page 3

BEST AVAILABLE COPY

ORNAMENTAL INSECT CONTROL

McNess Garden Dust can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, gladiolus, chrysanthemum, etc., a butterfly jumper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not apply to Creeping or Boston Ivy. Injury to tender foliage may occur if plants are wet when treated or in presence of high humidity.

Apply to the upper and lower leaf surfaces of all foliage for complete coverage to control such pests as:

Apple Aphids
Bagworms
Birch Leaf Miners
Blister Beetles
Roselard Bugs
Boxwood Leaf Miners

Elm Leaf Aphids
Elm Leaf Beetles
Fire Beetles
Gypsy Moths
Japanese Beetles
June Beetles

Lace Bugs
Leafhoppers
Mealy Bugs
Minute Webworms
Oak Leaf Miners


Orange Tortrix
Periodical Cicada
Plant Bugs
Pine Caterpillars
Psyllids
Rose Aphids

Rose-slug
Scale Insects
Tent Caterpillars
Thorn Bugs
Thrips (see page 4)
Willow Leaf

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are present in the treatment area.



ACTIVE INGREDIENTS:

O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 22.2%
Aromatic Petroleum Distillate 67.8%

AMENDMENT
To Previously Registered Labeling

Amend As Follows:

RECOMMENDED APPLICATIONS

CROP	INSECT	Pints GUTHION 2L	REMARKS
Cotton (Conven- tional or Low Volume Spray)	Boll weevil	1/2 to 1	Apply specified dosage per acre by air or ground equipment in at least 1 gal- lon of water. For early season control of bollworm and pink bollworm use 2 to 3 pints and for mid-to-late season applications use 3 to 4 pints. Repeat as necessary. Do not apply within 1 day of picking. Cotton receiving late season applications should not be pastured. <u>NOTE: GUTHION 2L at rates up to 1 pint per acre may be tank mixed with Galecron 4E or Fundal 4EC for ovicidal control of bollworms and tobacco budworms. For control of the larval stages of resistant and non-resistant bollworm and/or tobacco budworm, BOLSTAR 6 may be included in GUTHION 2L tank mixes. When using GUTHION 2L in tank mixes, observe all limitations, precautions and rate recommendations which appear on the labels for these products.</u>
	Brown cotton leafworm	1	
	Cotton fleahopper		
	Cotton leafworm		
	Lygus bugs		
	Thrips		
	Rapid plant bug	1 to 2	
Tarnished plant bug			
	Stink bug	2	
	Bollworm	2 to 4	
	Pink bollworm		

and
Fundal
4 EC
2a.

All copy except that which is underscored has been previously registered.