

GARDEN DUST

A Combination Fungicide and Insecticide
Contains Sevin® brand carbaryl insecticide
Ready To Use — Needs no mixing or diluting —
 simply use as it comes from the package.
Highly Effective — Kills listed insects and will
 control many of the most prominent destructive
 blights or fungus diseases.



A Dual Purpose Product — Highly effective against listed insects and
 fungi. It is no longer necessary to purchase several different products
 to give good control in your garden. You can rely on this product to
 safeguard your plants and shrubs effectively.

Economical to Use — Helps protect your investment in
 your flower and vegetable gardens and ornamental
 shrubs.

ACTIVE INGREDIENTS:

Maneb (Manganese ethylene bisdithiocarbamate)

Carbaryl (1 Naphthyl Methylcarbamate)

INERT INGREDIENTS: (including ZnSO₄)

34.5
 50.5
 95.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel For Additional
 Precautionary Statements

E.P.A. Reg. No. 2393-408

E.P.A. Est. No. 44616-MO-1

*Registered trademark of Union Carbide Corporation
 for the active ingredient carbaryl

NET WEIGHT 1 POUND



SOLD BY



Hopkins

agricultural chemical co.

Box 7532 • MADISON WISCONSIN 53702
 U.S.A.



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

HARMFUL IF INHALED OR SWALLOWED Do not breathe dust. Avoid contact with eyes and clothing. Do not take internally. Skin contact may be harmful. Avoid skin contact. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container only, preferably in a locked area. Keep container tightly closed when not in use. Store in a cool, dry place. High temperatures may shorten shelf life of product. Do not use food or drink containers for mixing or storage.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

Apply as soon as plants become infested with insects and damage is noticeable. Apply at 5 to 10 day intervals as needed. Dust thoroughly to get good coverage on upper and lower sides of leaves. Use a good dusting dust attractant buffer in late evening or early morning when vegetation is moist with dew and when the air is calm.

Subsequent planting of food and feed crops in treated soil must be made with crops on this product carbaryl label.

BEANS: - for Cowpeas, Black-eyed Peas, - for Downy Mildew, Lima, Snap, Green; - for Anthracnose, Early Mildew, Cowpeas, Black-eyed Peas, Lima, Snap, Green; - for Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Flea Beetles, Japanese Beetles, Leafhoppers, Velvetbean Caterpillars, Western Bean Cutworms, Armyworms, Corn Earworms, Stink Bugs, Tarnished Plant Bugs. Dust plants thoroughly at first sign of disease or infestation. Repeat at 5 to 10 day intervals. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

CARROTS: - for Cercospora Leaf Spot, Alternaria, Flea Beetles, Leafhoppers, Harlequin Bugs. Begin when plants are 6 weeks old. Repeat at 7 to 10 day intervals.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER: - for Alternaria Leaf Spot, Downy Mildew, Flea Beetles, Harlequin Bugs, Armyworms, Cabbage Worms and Loopers, Corn Earworms. Dust plants thoroughly at first sign of disease or infestation. Repeat at 3 to 7 day intervals. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

SWEET CORN: - for Helminthosporium Leaf Blight, Corn Earworms, Corn Rootworm Adults, European Corn Borers, Fall Armyworms, Flea Beetles, Japanese Beetles, Sap Beetles and Leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ear, apply at 3 day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Do not feed treated forage to dairy animals or to animals being finished for slaughter. Applications of this product to the tassels region of corn during the pollen shed period will seriously reduce bee populations.

MELONS: - for Downy Mildew, Anthracnose, Cercospora Leaf Spot, Burning Stem Blight, Flea Beetles, Melonworms, Flea Beetles, Leafhoppers. Begin at late bunching when vines begin to run or when disease or infestation first appears. Do not use on watermelons in Florida. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

CUCUMBERS: - Downy Mildew, Anthracnose, Alternaria (Macrosporium) Leaf Spot, Angular Leaf Spot, Phythium Fruit Rot, Pickleworms, Cucumber Beetles, Flea Beetles, Leafhoppers. Begin when plants begin to run or when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

PUMPKINS: - for Downy Mildew, Angular Leaf Spot, Flea Beetles, Leafhoppers, Melonworms. Begin when disease or infestation first begins. Repeat at 7 to 10 day intervals.

SQUASH: - For Downy Mildew, Anthracnose, Flea Beetles, Leafhoppers, Squash Bugs. Begin when plants start to run or when infestation first appears. Repeat at weekly intervals. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

GRAPE: - for Black Rot, Grape Berry Moths, Japanese Beetles, June Beetles, Red Banded Leaf Feeders, Grape Leaf Feeders, Grape Leafhopper, and Grape Leaf Skeletonizers. Begin just before bloom for Black Rot. Apply as necessary. Apply just before first brood Leaf Folder larvae emerge from rolls and as needed for Leafhoppers.

COLLARDS, KALE: - for Downy Mildew, Alternaria Leaf Spot, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

HEAD-LETTUCE: - Downy Mildew, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

SPINACH: - for Downy Mildew, White Rust, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

CRISPANT: - for Anthracnose, Early Blight, Alternaria, Phomopsis Blight or Fruit Rot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Leaf Bugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when fruit starts to form or when infestation appears. Repeat at 7 to 10 day intervals.

PEPPERS: - for Anthracnose, Cercospora Leaf Spot, Ripe Rot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals.

TOMATOES: - (Greenhouse and Field) for Cladosporium Leaf Mold, Early and Late Blight, Gray Leaf Spot, Stem Phythium, Septoria Leaf Spot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tomato Fruit Worms, Tomato Hornworms, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when fruit forms or infestation appears. Repeat at 7 to 10 day intervals. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.

DIRECTIONS FOR USE (Cont.)

- * ROSES — For Black Spot, Cercospora Leaf Spot, Rust. Begin when first leaves unfold. Repeat at 7 to 10 day intervals.
/ BOXELDER BUGS — Dust around foundation of house and a few feet of surrounding lawn.
GRASSHOPPERS on above crops. Dust as necessary.

ORNAMENTAL INSECT CONTROL

This product can be safely used on a wide range of ornamental plants, flowers, shrubs, and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arborvitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not apply to Virginia Creeper or Boston Ivy. Injury to tender foliage may occur if plants are wet when treated or in presence of high humidity.

Apply to the upper and lower leaf surfaces of all foliage for complete coverage to control such pests as:

Apple Aphids	Japanese Beetles	Puss Caterpillars
Bagworms	June Beetles	Psyllids
Birch Leaf Miners	Lace Bugs	Rose Aphids
Blister Beetles	Leafhoppers	Rose Slug
Boxelder Bugs	Mealy Bugs	Scale Insects
Boxwood Leaf Miners	Mimosa Webworms	Tent Caterpillars
Elm Leaf Aphids	Oak Leaf Miners	Thorn Bugs
Elm Leaf Beetles	Orange Tortrix	Thrips (exposed)
Flag Beetles	Periodical Cicada	Willow Leaf Beetles
Gypsy Moths	Plant Bugs	

NOTICE

Manufacturer and Seller warrant that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. Manufacturer and Seller make no other express or implied warranty of fitness or merchantability. The exclusive remedy of the buyer or user, and the limit of the liability of the Manufacturer and Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice, which may not be varied by any verbal or written agreement.