

STATEMENTS OF PRACTICAL TREATMENT

Call a physician or Poison Control Center immediately.

IF SWALLOWED: Induce vomiting by drinking 1 or 2 glasses of water and reaching back of the throat with finger. Use syrup of ipecac if available. Do not induce vomiting if person is unconscious. Do not give anything by mouth to an unconscious person.

IF IN EYES: Flush eyes with plenty of water without delay. Get medical attention.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Remove person to fresh air and give first aid as directed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed and absorbed through skin. Avoid breathing vapors of spray mist. Avoid contact with skin and eyes. In case of contact immediately flush eyes with plenty of water. Obtain medical attention if fatal symptoms.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in food processing plants, wash all equipment with potable water, benches, shelving, etc. where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds, and cover fish aquariums before spraying.

PHYSICAL HAZARD

Do not apply this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazards. If there is a need to treat these areas, use an oil base spray.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Avoid temperatures below 32°F.

PESTICIDE DISPOSAL: Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL: Triple rinse or equivalent and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by State and local authorities.

INDOORS

Avoid spotting of wallpaper, fabrics and other furnishings. Check materials for color fastness by dampening an inconspicuous area with this product and pressing a white tissue to the dampened area. If color transfer or change occurs, do not use this product.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: TO CONTROL ROACHES, ANTS, SILVERFISH, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under panels, around containers of stored food, around the base of machinery, behind shelves and

Additional Use Instructions on Right Panel


Hopkins

PESTRIN AQUA SPRAY

EFFECTIVE, FAST ACTING GENERAL PURPOSE INSECTICIDE

For Use In: Homes, Institutions, Restaurants, Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control.

Apply With Conventional Mechanical or Compressed Air Equipment.



ACTIVE INGREDIENTS:

Pyrethrins 0.1%
*Piperonyl Butoxide, Technical 1.0%
Petroleum Distillate 0.4%

INERT INGREDIENTS: 98.5%

TOTAL 100.0%

* Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panels For Additional Precautionary Statements



Hopkins

agricultural chemical co.

BOX 7532 • MADISON, WISCONSIN 53707
U.S.A.

SHAKE WELL BEFORE USE

NET CONTENTS

GALLONS

Hopkins

PESTRIN AQUA SPRAY

**EFFECTIVE, FAST ACTING
GENERAL PURPOSE INSECTICIDE**

For Use In: Homes, Institutions, Restaurants, Food Processing Plants,
Dairies, Livestock and Outdoors for Mosquito Control.
Apply With Conventional Mechanical or Compressed Air Equipment.



2393-1448

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	1.0%
Petroleum Distillate	0.4%
INERT INGREDIENTS:	98.5%

TOTAL

* Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl)
ether and 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panels For Additional Precautionary Statements



Hopkins
agricultural chemical co.

BOX 7532 • MADISON, WISCONSIN 53717
U.S.A.

SHAKE WELL BEFORE USE

NET CONTENTS

GALLONS

DIRECTIONS (Continued)

For use in homes, institutions, restaurants, food processing plants, dairies, livestock and outdoors for mosquito control. Apply with conventional mechanical or compressed air equipment. Repeat treatment as needed.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make a second application of floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray headboards. Treat baseboards, mouldings and floors. Repeat treatment as needed.

TO CONTROL FLEAS AND BROWN DOG TICKS: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage areas directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon per 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

LIVESTOCK

To protect cattle, beef, dairy and horses (including food animals) from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

When using this product on livestock, care should be taken that the spray does not come in direct contact with the lactating dairy cow's teats unless they are washed with an antiseptic and cleansing solution and dried before milking.

OUTDOORS

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools, ponds and shore lines. Apply at a rate of 1 gallon per acre. 4-6 ounces per 1000 square feet. Use a mist blower or other conventional misting equipment.

Using truck-mounted conventional misting equipment traveling at 5 mph, calculate equipment to deliver 4-6 gallons per minute to treat a 300 foot swath.

NOTICE OF WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material. After purchase and handling is contrary to label instructions. Buyer agrees in purchasing this product that in the event of damages arising from a breach of the warranty, he will accept a refund of the purchase price of the product, less the cost of the manufacturer's label.

*Subject to applicable state laws.

EPA Reg No 2393-448

EPA Est No 2393-IL-3 • 2393-WI-1 •

523-IL-1 • 2393-WI-2 • 33509-MO-1 •

Superscript refers to first letter of lot number