

### DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**FOR INDOOR FOGGING:** Adjust nozzle to position for a very fine mist. Close all openings, air conditioning and ventilating systems. Cover all foodstuffs to prevent contamination from falling insects. Sweep up and destroy all dead insects. Use 1 gallon of concentrate for each 50,000 cubic feet of space, or wet surfaces thoroughly and treat all surface areas, insect hiding places and runways at the rate of 1 gallon to 750 square feet Repeat when infestation occurs.

**FOR FOOD PROCESSORS:** Cover or remove all food before applying above treatments. Thoroughly clean all food processing surfaces before reusing Apply only when plant is not in operation. Do not remain in treated areas Ventilate the area after treatment is completed Remove pets, birds and cover fish aquariums before spraying

### CAUTION


**FLAMMABLE** - Do not spray into or near open flame. Do not smoke while using Keep container away from stoves, radiators and all other hot surfaces May be harmful if swallowed, inhaled or absorbed through the skin Avoid breathing of spray mist and provide adequate ventilation of area being treated Contact with skin, eyes or clothing should also be avoided. Wash thoroughly after using Avoid contamination of food, utensils and food preparation areas

# FOGGING SPRAY CONCENTRATE

FOR CONTROL OF GRANARY WEEVIL, RICE WEEVIL, CONFUSED FLOUR BEETLE, RUST RED FLOUR BEETLE, SAW-TOOTHED GRAIN BEETLE, SPIDER BEETLE, ANGOUMOIS GRAIN MOTH and MEDITERRANEAN FLOUR MOTH.

### ACTIVE INGREDIENTS

- Petroleum Distillate 99.02%
- N Octyl Bicycloheptene Dicarboximide 0.52%
- \* Technical Piperonyl Butoxide 0.31%
- Pyrethrins 0.15%
- \* Equivalent to 0.25% (ibutylcarbityl) (6-propylpiperonyl) ether and 0.06% related compounds. 100.00%



**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
See Right Side Panel For First Aid Statements.

**FOR COMMERCIAL AND INDUSTRIAL USE ONLY, OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS.**

MANUFACTURED BY  
**I. SCHNEID, INC.**  
1429 FAIRMONT AVE. NW  
ATLANTA GA 30381-4601

### PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock.

### STORAGE AND DISPOSAL

**STORAGE:** Keep container closed when not in use.  
**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Do not contaminate water, food or feed by storage or disposal.  
**CONTAINER DISPOSAL:** Triple rinse (or equivalent) Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities

### FIRST AID

In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, promptly drink a large quantity of water. Avoid alcohol. Call a physician immediately.

**CONTROL NUMBER**

**NET CONTENTS GALLONS**

**ENVIRONMENTAL HAZARD:**

Keep Out of Lakes,  
Streams or Ponds.

**DIRECTIONS FOR USE**

It is a violation of federal law to use  
this product in a manner inconsistent with  
its labeling.

**DISPOSAL**

Do not reuse empty container.  
Wrap container and put in trash.

**USE RESTRICTIONS** - For control of Norway  
rats, Roof rats and House mice in and  
around homes, industrial and agricultural  
buildings, and similar man-made structures.

Treated baits must be placed in tamper-  
proof bait boxes or in locations not  
accessible to children, pets, domestic  
animals or wildlife. Do not place bait in  
areas where there is a possibility of  
contaminating food or surfaces that come in  
direct contact with food.



**ACTIVE INGREDIENT**

Diphacinone [2-(Diphenylacetyl)-  
1,3-indandione] . . . . . 0.005%

**INERT INGREDIENTS** . . . . . 99.995%

**TOTAL** . . . . . 100.000%

EPA Reg. No. 2393-476

EPA Est. No. 2393-WI-1

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See Other Panel For Additional Precautions

MANUFACTURED BY

*Hopkins* agricultural chemical co.  
BOX 7532 • MADISON, WISCONSIN 53707 U.S.A.

NET WEIGHT 1 LB.

supply of fresh bait for at least 10 days.

**MICE:** Apply 1/4 to 1/2 ounce (1-2 level  
tablespoons) of bait at 8 to 12 foot  
intervals. Larger placements (up to 2  
ounces) may be needed at points of very  
high mouse activity. Maintain an uninter-  
rupted supply of fresh bait for at least 15  
days.

**RATS & MICE:** Replace contaminated or  
spoiled bait immediately. Collect and  
dispose of all dead animals and unconsumed  
bait properly. To prevent reinfestation,  
eliminate food, water, and harborage as  
much as possible. If reinfestation does  
occur, repeat treatment. Where a con-  
tinuous source of infestation is present,  
establish permanent bait stations and  
replenish bait as needed.

**NOTICE OF WARRANTY**

SELLER AND MANUFACTURER MAKE NO WARRANTY  
OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE,  
EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT  
OR ITS USE, WHICH EXTEND BEYOND THE STATE-  
MENTS ON THIS LABEL.

**ACCEPTED**

MAR 3 1986

**BEST AVAILABLE COPY**

U-  
7-  
of  
now  
EPA Reg. No.

2393-476

# RAT AND MOUSE KILLER

WITH DIPHACINONE

A Pelletized Rodenticide

READY MIXED, EASY TO USE — USE INDOORS AND  
OUTDOORS



#### ACTIVE INGREDIENT

Diphacinone [2-(Diphenylacetyl)-  
1,3-indandione] . . . . . 0.005%  
INERT INGREDIENTS . . . . . 99.995%

TOTAL . . . . . 100.000%

EPA Reg. No. 2393-476

EPA Est. No. 2393-WI-1

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

See Other Panel For Additional Precautions

MANUFACTURED BY

*Hopkins* agricultural chemical co.  
BOX 7532 • MADISON, WISCONSIN 53707 U.S.A.

NET WEIGHT 1 LB.

#### PRECAUTIONARY STATEMENTS:

##### HAZARD TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

**NOTE TO PHYSICIAN -** If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in Dihydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

##### ENVIRONMENTAL HAZARD:

Keep Out of Lakes,  
Streams or Ponds.

##### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### DISPOSAL

Do not reuse empty container.  
Wrap container and put in trash.

**USE RESTRICTIONS -** For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS -** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

#### APPLICATION DIRECTIONS

**RATS:** Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

#### NOTICE OF WARRANTY

SELLER AND MANUFACTURER MAKE NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL

ACCEPTED

MAR 3 1966

2393-476

PAI 16  
2092-4/16  
1.0.0