ACID SANITIZER

• FOR THE DAIRY INDUSTRY• CLEANS • REMOVES MILKSTONE • SANITIZES

DIRECTIONS

Elko Acid Sanitizer is easily dispersed in water up to 1700 ppm of hardness to form a completely uniform solution. When used at 1:500 aqueous dilution (200 ppm active quarternaries) may be used on food processing and dairy equipment without need of final potable water rinse.

CLEANING BULK MILK TANKS: As soon as possible after milk has been removed from tank, rinse completely with cold water and drain. Place 4 gal. warm water (120°F) in tank and add 1 ounce (2 Table spoons) Elko Acid Sanitizer. Scrub lank, mixers and cover and drain. SANITIZING MILK EQUIPMENT AND COW UDDERS: Previously cleaned milking equipment, pails, inflations, and tubing should be sanitized prior to milking. Mix Elko Acid Sanitizer as above, scrub or dip equipment, drain before beginning milking operation.

Use clean single service towel for each animal. Wipe udders with above solution prior to milking. Never reuse towel or dip it back into * the solution. Always use freshly prepared solution and non-corrodible utensils.

ACTIVE INGREDIENTS:

- D A N G E R KEEP OUT OF REACH OF CHILDREN

See right side panel for First Aid and additional pred



ELKHORN CHEMICAL CO One Elko Lane Elkhorn, Wisconsin 53121

NET CONTENTS: ONE U.S. GA

000

0 0 0 0 0 0

IGREDIENTS:

C₁₄, 30 C₁₆, 5 / C₁₂, 5 (₁₈) benzyl ammonium chlorides

 $\sim C_{12}/17 + C_{14}$) dimethyl yl ammonton chlofides

GREDIENTS

OUT OF REACH OF CHILDREN

nel for First Aid and additional precautions

ELKHORN CHEMICAL CO., INC. One Elko Lane

Elkhorn, Wisconsin 53121

NET CONTENTS: ONE U.S. GAL.

- D A-N G E R -

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

Rinse empty container thoroughly with water and discard it.

- F | R.S T A 1 D -

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

PHYSICIAN - N O T E T 0

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Ll'A Reg. No. 2437-16

EPA LS1 #2431 WS-1

ACCEPTED

5.0%

30.0 % 60.0%