

Reduce to 47% strength

ACCEPTED

DEC 4 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 2439-8 SUBJECT
TO ATTACHED COMMENTS.

Apperson algaecide is efficient, economical and stable in use dilution.

Apperson is compatible with all chemicals used in pool water and will not damage tile, concrete, metal, or plastics.

Apperson keeps pool water free and sparkling, clear of visible algae, slime and green brown colors.

Apperson improves filter operation and cuts down need for other chemicals.

Apperson imparts tone and a pleasant "feel" to swimming pool water.

Do not reuse container. Destroy when empty.

This product is toxic to fish. Do not discharge pool water where it will drain into lakes, streams, ponds or public waters.

Apply this product only as specified on this label.

DANGER: Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

EPA REG. No. 2439-8

APPERSON

SWIMMING POOL

ALGAECIDE

OUTSTANDING ALGAE GROWTH DEPRESSANT

ACTIVE INGREDIENT:

Alkyl* Dimethyl Benzyl Ammonium Chloride
*(C₁₀, 60%; C₁₂, 25%; C₁₆, 15%) 10%

INERT INGREDIENTS 90%

"DANGER" 100%

"Keep Out of Reach of Children"
See other precautions on the
back side panel.

Manufactured By:

APPERSON CHEMICALS, INC.

JACKSONVILLE, FLORIDA

ONE GALLON

DIRECTIONS

Initial Application: To a cleaned pool add one gallon to 50,000 gallons of water, or ratio thereof.

Maintenance Dose: Use one quart per 50,000 gallons of water every 5 to 7 days. When weather is hot or bathing load is heavy, use this dosage every 5 days or as necessary.

Booster Dose: Immediately after every rain add one quart per 50,000 gallons of water as above.

Important: Follow above directions without fail, even when pool is not used. Algae growth must be fought at all times.

Pool-side surfaces: After cleaning surfaces, use one oz. to four gallons of water for sanitizing or one oz. to two gallons of water for disinfecting.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.