


Contains 6 Lbs.

*Toxaphene Per
Gallon



Master
B R A N D

Contains 6 Lbs.

*Toxaphene Per
Gallon

EMULSIFIABLE SPRAY FOR COTTON

SHAKE WELL BEFORE USING
CONTENTS BY WEIGHT

ACTIVE INGREDIENT:

*Toxaphene

58.5 Per Cent


INERT INGREDIENTS

41.5 Per Cent

TOTAL

100.0 Per Cent

*Technical Chlorinated Camphene
(Chlorine Content 67 per cent—69 per cent)



DANGER

KEEP OUT OF THE REACH OF CHILDREN

SEE OTHER PANEL FOR ADDITIONAL WARNINGS

WARNING: May be fatal if swallowed! May be absorbed through the skin! May produce irritation of skin and eyes. Avoid breathing of vapor or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed or foodstuffs.

Highly toxic to fish and wildlife, do not contaminate any body of water or apply under conditions favoring drift outside target area. Birds feeding on treated area may be killed.

This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

CAUTION: Do not apply to cucurbits. Do not apply to feed and forage crops if they are to be fed to dairy or beef animals being finished for slaughter. Do not apply to other crops unless specifically recommended by your State Agricultural Experiment Station.

COTTON: Bollweevil, Cotton Fleahoppers, Lygus Plant Bug, Thrips, Armyworms and Grasshoppers.

SPRAY EQUIPMENT

For effective control against cotton, MASTER SPRAY should be used with sprayers capable of delivering a certain amount of dilute spray per acre and of dilute spray used per acre will depend upon the type, make and speed of equipment and the number of nozzles used per acre. It is necessary to determine the nozzle output, nozzle pressure of at least 15 to 40 pounds per square inch, depending upon the nozzle size, and to maintain at least 10 inches from the plants to prevent burning of foliage.

RATE OF APPLICATION

MASTER SPRAY should be used at the recommended rate per acre mixed with the amount of water necessary to cover one acre. The gallonage of spray applied will vary from 1 to 10 gallons. Tractor mounted and self propelled sprayers will usually put out from 1 to 3 gallons of spray per acre. Airplane applications should be made at the rate of 1 to 2 gallons per acre.

NOZZLE ARRANGEMENT FOR GROUND EQUIPMENT

SEEDLING COTTON: Use one nozzle per row placed 10 inches directly above the plants.

COTTON UP TO 20 INCHES HIGH: Use two nozzles per row placed one on each side of the row and 10 inches from the plant. The nozzles should be directed so that they cover the top and sides of each plant.

COTTON OVER 20 INCHES HIGH: Use three nozzles per row: one placed 10 inches directly above the plant and one on each side.

COTTON OVER 20 INCHES HIGH: Use three nozzles per row: one placed 10 inches directly above the plant and one on each side.

CONTROL RECOMMENDATIONS

Recommendations vary in the different cotton producing areas and it is suggested that locally issued state recommendations be followed for proper timing and number of applications to be made. Infestation counts should be made in each field by the grower and applications made as necessary. Spraying can be done over a wide range of conditions almost at any time of the day. It should not be done during rain or when the wind is excessively high.

THRIPS, COTTON FLEAHOPPERS, LYGUS PLANT BUGS CONTROL ON SEEDLING COTTON: Use 1 quart of MASTER SPRAY per acre.

EARLY SEASON BOLLWEEVIL CONTROL: Use 1/3 gallon of MASTER SPRAY per acre. Make 2 to 3 applications at 7 to 10 day intervals.

MIDSEASON CONTROL: Use 3 parts of MASTER SPRAY per acre. Make the first application when infestation counts indicate its necessity. Two or three additional applications should be made at 5 day intervals until the infestation is brought under control. Additional applications may be necessary.

LATE SEASON CONTROL: Use 3 to 4 parts of MASTER SPRAY per acre. Make applications at 5 day intervals as necessary. If control is necessary after ammunition begins, the applications should be made at intervals of 3 days or less.

ARMYWORM AND FALL ARMYWORM (GRASSWORM): Use at the rate of 1/3 gallon per acre diluted with the volume of water applied by the sprayer being used.

GRASSHOPPERS: Use at the rate of 1/3 gallon per acre diluted with the volume of water applied by the sprayer.

NOTICE

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated in the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Stevens Industries, Inc. and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.

Manufactured By

STEVENS INDUSTRIES, INC.

DAWSON, GEORGIA 31742

U.S.D.A. Reg. No. 2459-30