

INSECT ON PASTURE BRAND PRODUCTS

DIRECTIONS

USE:

FOR THE CONTROL OF:


GRASSHOPPERS
ARMY WORMS
THrips
HORNWORMS
STINK BUG

VEGET BROWN CATERPILLAR
LEAF HOPPERS
CORNEA CURCULIO
BLISTER BEETLES
FLEA HOPPERS

ATTACKING:

PEAS
BRUSSELS SPROUTS
PEPPERS
ONIONS

BROCCOLI
CABBAGE
EGGPLANT
CORN



Dust upper and lower leaf surface when air is still. Repeat as necessary. Do not use on green or spring onions. Do not use on peppers, melons, squash, eggplant after fruit starts to form. Do not use on other crops listed within ten days of harvest. Do not apply to cabbage after head starts to form or 30 days of harvest. Remove all residue at harvest by washing or other effective means. For commercial growers, dust at the rate of 20 lbs. per acre.

Do not feed treated forage to dairy animals or to animals being finished for slaughter.

FOR LAWNS:

FOR THE CONTROL OF:

Ticks
Cutworms

Fall Armyworms
Chiggers

Apply at the rate of one pound per 2000 square feet (20 lbs. per acre). Birds feeding on treated areas may be harmed.

CAUTION

BRUSSELS SPROUTS
PEPPERS
ONIONS

CABBAGE
EGGPLANT
CORN

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 2459-119

Dust upper and lower leaf surface when air is still. Repeat as necessary. Do not use on green or spring onions. Do not use on peppers, melons, squash, eggplant after fruit starts to form. Do not use on other crops listed within ten days of harvest. Do not apply to cabbage after head starts to form or 30 days of harvest. Remove all residue at harvest by washing or other effective means. For commercial growers, dust at the rate of 20 lbs. per acre.

Do not feed treated forage to dairy animals or to animals being finished for slaughter.

FOR LAWNS:

FOR THE CONTROL OF:

Ticks
Cutworms

Fall Armyworms
Chiggers

Apply at the rate of one pound per 2000 square feet (20 lbs. per acre). Birds feeding on treated areas may be harmed.

C A U T I O N

KEEP CHILDREN AND PETS OFF THE TREATED AREAS UNTIL THE INSECTICIDE HAS BEEN WASHED INTO THE SOIL AND THE TREATED AREAS HAVE DRIED COMPLETELY. HARMFUL IF SWALLOWED! AVOID PROLONGED BREATHING OF DUST. AVOID CONTACT WITH EYES, SKIN AND CLOTHING -- WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

Do not apply when weather conditions favor drift of dust from crop area treated. Do not feed treated crops or crop refuse to dairy animals or animals being finished for slaughter. Do not allow grazing of crops treated with this material. Do not apply to varieties of peas with edible pods after the pods start forming. Do not use on Broccoli and Brussels Sprouts after the edible parts start to form.

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be harmed.