

This	ACTIVE INGREDIENT	
Package	Carbaryl (1-Naphthyl N-Methylocarbamate)	5.00%
Contains:	INERT INGREDIENTS	95.00%
	Total	100.00%

*Sevin is a trade mark of Union Carbide Corp. for Carbaryl (1-Naphthyl N-Methylocarbamate).

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles.

DIRECTIONS—Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary for good control. Read entire label. Use in strict accordance with label directions and cautions.

VEGETABLES—Beans: Use 10 pounds per acre for Mexican bean beetle, 20 pounds per acre (1 ounce per 50' row) for bean leaf beetle, cucumber beetles, flea beetles, leafhoppers, velvet bean caterpillar, armyworms, corn earworm, stink bugs and tarnished plant bug. This product can be applied up to the date of harvest.

Corn: Use 20 to 40 pounds per acre (1.2 ounces per 50' row) for corn earworm, European corn borer, fall armyworm, flea beetles and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders apply to entire plant. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Do not apply this product to corn within seven days of harvest for forage.

Potato, tomato, pepper and eggplant: Use 10 to 20 pounds per acre for Colorado potato beetle, flea beetle, and leafhoppers. Use 20 to 40 pounds for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. This product can be applied up to the date of harvest. Okra, cucumber and squash: Use 20 pounds per acre for cucumber beetles, flea beetles, leafhoppers, squash bugs, corn earworm and stink bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

POULTRY—On chickens, turkeys, ducks, partridges, pheasants and pigeons: For Northern fowl mite and lice, as a supplement to premise treatment for chicken mite and fleas: Use 1 pound per 100 birds. Apply thoroughly by means of a shaker can or hand duster. Do not apply to poultry within seven days of slaughter. Do not repeat applications within four weeks. Do not contaminate food and drinking water.


Poultry Houses: For chicken mite, fleas and bedbugs. Apply 1 pound per 40 square feet of litter or floor space, and to roosts and adjacent areas by means of a hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not house birds in treated houses within seven days of slaughter. Do not contaminate feed and drinking water.

PETS—On dogs and cats. For ticks and fleas dust liberally over the entire animal. Rub thoroughly into skin. Dust in and around sleeping quarters and other areas where ticks and fleas occur. Repeat weekly if needed. Do not treat kittens less than 4 weeks old.

ORNAMENTALS—To control bagworm, leaf miner, blister beetles, elm leaf aphid, flea beetles, Japanese beetles, lace bugs, leaf rollers, plant bugs apply thoroughly to the infested plants at the first sign of damage and repeat at 7 day intervals or as necessary for good insect control. To obtain scale control apply in spring and early summer when scale crawlers are present.

NOTICE

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express, or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to Stevens Industries, Inc., and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.



CONTENTS

**5%
SEVIN**

CAUTION

**KEEP OUT OF THE REACH OF
SEE OTHER PANEL FOR ADDITION**

This	ACTIVE INGREDIENT
Package	Carbaryl (1-Naphthyl N-Methylocarbamate)
Contains:	INERT INGREDIENTS

Total ...

*Sevin is a trade mark of Union Carbide Corp. for N-Methylocarbamate).

U.S.

**Manufactured by
STEVENS INDUSTRIES, INC., D**

(Methylocarbamate) 5.00%
 95.00%
 Total 100.00%
 Corp. for Carbaryl (1-Naphthyl)

Take internally. Avoid breathing
 Wash thoroughly after handling.
 Avoid contamination of foodstuffs. Avoid con-
 tact with eyes.

coverage. Apply when insects or
 or as necessary for good control.
 Follow directions and cautions.

Mexican bean beetle, 20 pounds
 cucumber beetles, flea beetles
 corn earworm, stink bugs and
 to the date of harvest.

per 50' row for corn earworm,
 and leafhoppers. For larvae in
 apply to entire plant. For insects
 starting when first silks appear
 no applications may be required
 good coverage are absolutely es-
 sential to corn within seven days of

10 pounds per acre for Colorado
 to 40 pounds for European corn
 n, tomato hornworm. Tarnished
 applied up to the date of harvest
 acre for cucumber beetles, flea
 and stink bugs. Avoid excessive
 treatments are made when tender foliage

chickens, pheasants and pigeons: For
 mite treatment for chicken mite
 apply by means of a shaker can or
 days of slaughter. Do not repeat
 feed and drinking water.

Apply 1 pound per 40 square feet
 as by means of a hand duster or
 repeat as necessary. Do not treat
 of house birds in treated houses
 feed and drinking water.

liberally over the entire animal.
 quarters and other areas where
 Do not treat kittens less than 4

miner, blister beetles, elm leaf
 leaf rollers, plant bugs apply
 of damage and repeat at 7 day
 obtain scale control apply in spring

This product conforms to its chemi-
 cal purpose stated on the label
 under normal conditions of use,
 warranty of MERCHANTABILITY-
 express, or implied, extends
 instructions, or under abnor-
 mally foreseeable to Stevens
 of any such use. Any dam-
 age shall be limited to direct
 actual commercial damage such



CONTENTS

1 POUND

5% SEVIN*

CAUTION

KEEP OUT OF THE REACH OF CHILDREN
SEE OTHER PANEL FOR ADDITIONAL WARNINGS

This	ACTIVE INGREDIENT	
Package	Carbaryl (1-Naphthyl N-Methylcarbamate)	5.00%
Contains:	INERT INGREDIENTS	95.00%
	Total	100.00%

*Sevin is a trade mark of Union Carbide Corp. for Carbaryl (1-Naphthyl N-Methylcarbamate).

U.S.D.A. Reg. No. 2459-134

Manufactured by
STEVENS INDUSTRIES, INC., DAWSON, GEORGIA

31742

