

2459-165

See Antidote Statement and Other Required Warning Statements on the Other Panels.



Master
BRAND

NET CONTENTS
GALLON

VAPONA[®] INSECTICIDE EMULSIBLE CONCENTRATE

FOR USE BY PEST CONTROL OPERATORS ONLY.

Do not reuse this container; Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

DANGER

KEEP OUT OF THE REACH OF CHILDREN.

POISON



Do not use, pour, spill, or store near heat or open flame.

(Containing 2.0 lbs. VAPONA Insecticide per gal.)

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl dimethyl phosphate
Related Compounds
Petroleum Hydrocarbons

INERT INGREDIENTS

*Equivalent to 22.5 per cent w VAPONA Insecticide.

By Weight

22.0 Per Cent*
1.7 Per Cent*
71.3 Per Cent
5.0 Per Cent

TOTAL

100.0 Per Cent

VAPONA Emulsifiable and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See antidote and warnings below)

Store the emulsifiable in a safe place. Children or animals must not be able to reach or handle the emulsifiable or diluted emulsion. It is recommended that only the amount of diluted emulsion that will be used immediately be prepared from the emulsifiable.

Mix the emulsifiable into water outdoors or in a well-ventilated room. Wash off the outside of the bottle of emulsifiable in running water before storing away.

Manufactured by
STEVENS INDUSTRIES, INC.
DAWSON, GEORGIA

EPA Reg No 2459-165

EPA Est No 2459-Ga-1

This product is toxic to fish, birds, and other wildlife: Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

EXCEPT FOR CRACK AND CREVICE TREATMENT, DO NOT USE IN THE EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERCIALY PREPARED OR PROCESSED. DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSED.

DANGER

VAPONITE 2 Emulsifiable is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes; Avoid exposure to fumes; Wear clean rubber gloves, waterproof protective clothing and goggles when handling the emulsifiable to prepare emulsions.

Wear a mask or respirator of a type passed by the US Bureau of mines for DDVP (VAPONA Insecticide) protection. Keep all unprotected persons out of operational area!

In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water before eating or smoking; Wash all contaminated clothing with soap and hot water before reuse.

DECONTAMINATE AND DESTROY USED CONTAINERS:

Decontaminate used containers by washing thoroughly with a dilute solution of caustic (less than 5%) and detergent. **CAUTION:** Avoid skin and eye contact with caustic solutions! Rinse containers with water and dispose of the washings in such a way that you do not constitute a hazard to, or contaminate water supplies. Destroy containers by crushing and puncturing; then bury or burn; if burned, stay out of smoke

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR VAPONA POISONING; 2-PAM is also antidotal and may be used in conjunction with atropine! **CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING!**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt in a glass of water); **NEVER GIVE ANYTHING BY MOUTH TO A UNCONCIOUS PERSON!**

EXTERNAL: If the material has been spilled on skin, immediately remove patient from the vicinity of the VAPONA, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 10 minutes. **IF WARNING SYMPTOMS APPEAR: NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR!** (See WARNING SYMPTOMS below) Keep patient prone and quiet; Start artificial respiration immediately if patient is not breathing; Transport the patient immediately to the nearest hospital.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against para-sympathetic nerve stimulation. If there are signs of para-sympathetic stimulation, atropine sulfate should be injected at ten-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Pralidoxime chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining atropinization: In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion of 250 cc of saline over a 15 to 30 minute period; If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5% solution in water over not less than two minutes; After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved.

For info chest b tinuous cause further

TO BE OUTDOCK DOCKS, TIONS,

Apply V ants, sov To prep Apply as humans.

MANUR: apply at DOG KE

dog runv For the 1000 squ CAUTIO

HOUSEI mosquito contain

To prep For Hou pressure boards, areas w until sur

APP Places of served

LIMITI vering a. and crev bases and spiders b introduci

APPLIC/ MENTS,

Stevens I reasonabl normal co ABILITY product c reasonabl Any dam shall not

2459-165

This product is toxic to fish, birds, and other wildlife: Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

EXCEPT FOR CRACK AND CREVICE TREATMENT, DO NOT USE IN THE EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERCIALY PREPARED OR PROCESSED. DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSED.

DANGER

VAPONITE 2 Emulsifiable is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes; Avoid exposure to fumes; Wear clean rubber gloves, waterproof protective clothing and goggles when handling the emulsifiable to prepare emulsions.

Wear a mask or respirator of a type passed by the US Bureau of mines for DDVP (VAPONA Insecticide) protection. Keep all unprotected persons out of operational area!

In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water before eating or smoking; Wash all contaminated clothing with soap and hot water before reuse.

DECONTAMINATE AND DESTROY USED CONTAINERS:

Decontaminate used containers by washing thoroughly with a dilute solution of caustic (less than 5%) and detergent. CAUTION: Avoid skin and eye contact with caustic solutions! Rinse containers with water and dispose of the washings in such a way that you do not constitute a hazard to, or contaminate water supplies. Destroy containers by crushing and puncturing; then bury or burn; if burned, stay out of smoke

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR VAPONA POISONING: 2-PAM is also antidotal and may be used in conjunction with atropine! CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING!

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt in a glass of water): NEVER GIVE ANYTHING BY MOUTH TO A UNCONCIOUS PERSON!

EXTERNAL: If the material has been spilled on skin, immediately remove patient from the vicinity of the VAPONA, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 10 minutes. IF WARNING SYMPTOMS APPEAR: NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR! (See WARNING SYMPTOMS below) Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against para-sympathetic nerve stimulation. If there are signs of para-sympathetic stimulation, atropine sulfate should be injected at ten-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Pralidoxime chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining atropinization; In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion of 250 cc of saline over a 15 to 30 minute period; If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5% solution in water over not less than two minutes; After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved.

For infants and children, the dose of 2-PAM is 0.25 grams; Morphine is contra-indicated; Clear chest by postural drainage; Oxygen administration may be necessary; Observe patient continuously for 48 hours; repeat exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor; Allow no further exposure until cholinesterase regeneration has occurred as determined by blood test.

DIRECTIONS FOR USE

TO BE APPLIED BY PEST CONTROL OPERATORS OR TRAINED PERSONNEL ONLY
OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREA, PATIOS, LOADING DOCKS, OUTDOOR LATRINES, PARKING AREA, REFUSE AREAS AROUND SERVICE STATIONS, GARBAGE COLLECTION AND DISPOSAL AREAS

Apply VAPONA Insecticide as a 0.5% spray to control flies, gnats, mosquitoes, roaches, flea, ants, sowbugs, and brown dog ticks.

To prepare a 0.5% spray, dilute 1 pint of VAPONITE 2 Emulsifiable with 6 gallons of water; Apply as a coarse wet spray at the rate of 1 quart per 1000 square feet. Do not treat animals or humans. Repeat as necessary.

MANURE PILES, GARBAGE DUMPS: For the control of fly maggots in maggot-breeding areas, apply at the rate of 1-2 quarts of 0.5% spray per 1000 square feet.

DOG KENNELS: For the control of flies, mosquitoes, roaches, fleas and ants: Apply to outside dog runways window frames and ledges at a rate of 1 quart of 0.5% spray per 1000 square feet. For the control of fleas and brown dog ticks, increase dosage to 2 quarts of 0.5% spray per 1000 square feet.

CAUTION: Do not use this product in conduits, motor housings and electrical switch boxes.

HOUSEHOLD PEST CONTROL: Apply VAPONA Insecticide as a 0.5% spray to control flies, mosquitoes, wasp, gnats, small flying moths, cockroach, ant, spider and silverfish. Do not contaminate food, water, food containers and cooking utensils.

To prepare a 0.5% spray, dilute 1 pint of VAPONITE 2 Emulsifiable with 6 gallons of water; For Households, Theaters, Industrial Plants, Non-Food Warehouses, apply 0.5% spray with low pressure sprayer to localized areas insects may infest. Treat infested area around baseboards, cracks, walls, door and window frames, and localized areas of floors. Do not apply to areas which will be contacted frequently by children and do not allow them in treated areas until surfaces are dry. Do not treat animals.

APPLICATION WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

Places other than private residences in which exposed food is held, processed, prepared or served

LIMITED TO CRACK AND CREVICE TREATMENT ONLY (Using equipment capable of delivering a pin stream of insecticide) Apply a small amount of a 0.5% spray directly into cracks and crevices such as expansion joints between elements of construction or between equipment bases and the floor, wall voids, or hollow equipment legs where cockroaches, ants, silverfish or spiders hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATION OF THIS PRODUCT IN THE AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED

NOTICE

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Stevens Industries, Inc. and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.