

NET WEIGHT 50 LBS.



**Master**®

B R A N D

**5% MALATHION DUST**

RECEIVED  
MAY 12 1975

Under the Federal Insecticide, Fungicide, and Plant Disease Act,  
registered under Reg. No. 2459-181

THIS PACKAGE CONTAINS

ACTIVE INGREDIENT:

Malathion\* ..... 5%

INERT INGREDIENTS: ..... 95%

TOTAL 100%

\*O, O-Dimethyl Dithiophosphate of diethyl mercaptosuccinate

#### RECOMMENDATIONS

**COTTON:** For the control of Boll Weevil, Cotton Aphid, Cotton Leafworm, Spider Mites and Thrips apply 20 pounds per acre. Consult your State Agricultural Experiment Station or State Agricultural Extension Service for time and number of applications.

**TRUCK CROPS.** As indicated, apply at five to seven day intervals as necessary for control. Do not apply to plants unless they are dry as injury may occur. Dust at the rate of 35 lbs. per acre, equal to 1-3/4 oz. per 50 feet of row.

**BEANS—**For the control of Mexican bean beetles, leafhoppers, spider mites, Japanese beetles, aphids, cucumber beetles. Do not harvest within one day of application: Apply 30 to 35 lbs. per acre.

**COLE CROPS & LEAFY VEGETABLES—**Broccoli, Brussels sprouts, cabbage, collards, kale, mustard greens, parsley, turnips. For the control of aphids. Do not harvest within 7 days of application. Apply 30 to 35 pounds per acre.

**CUCURBITS—**Cucumber, melons. Do not apply within one day of harvest. For the control of aphids, pickle worms, leafhoppers, apply 32 to 35 lbs per acre.

**LETTUCE AND SPINACH—**For the control of aphids: Do not apply to head lettuce within 7 days of harvest or leaf lettuce within 14 days of harvest.

**PEAS—**For the control of pea aphids, pea weevils: Do not apply within 7 days of harvest. Apply 30 to 35 lbs per acre.

**POTATOES—**For the control of aphids, leafhoppers, mealy bugs. Apply 30 to 35 lbs per acre.

**TOMATOES—**For the control of aphids. Do not apply within one day of harvest. Apply 35 pounds per acre.

**POULTRY HOUSES—**For the control of Northern Fowl Mites, Chicken Red Mite and Poultry Lice, apply 3/4 pound per 50 to 60 square feet of litter, floor space, nests, roosts and adjacent areas. Repeat as necessary.

**CATTLE—**For the control of lice on beef cattle make a thorough application to the animals. Repeat application within 10 days and thereafter as needed. Avoid contamination of feed, food containers and watering troughs. Do not treat lactating dairy animals. Do not apply later than 5 hours before milking or during milking. Do not treat calves under one month of age.

**SWINE—**For the control of lice on swine, make a thorough application of the animals. Pens should be dusted: Repeat application in 10 days and thereafter as needed: Do not treat animals under 1 month of age.

#### CAUTION

KEEP OUT OF THE REACH OF CHILDREN

AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. HARMFUL IF SWALLOWED.

NOTE TO PESTICIDALISTS: Malathion may cause cholinesterase inhibition. Atropine is antidote.

\*O, O-Dimethyl Dithiophosphate of diethyl mercaptosuccinate

## RECOMMENDATIONS

**COTTON:** For the control of Boll Weevil, Cotton Aphid, Cotton Leafworm, Spider Mites and Thrips apply 20 pounds per acre. Consult your State Agricultural Experiment Station or State Agricultural Extension Service for time and number of applications.

**TRUCK CROPS.** As indicated, apply at five to seven day intervals as necessary for control. Do not apply to plants unless they are dry as injury may occur. Dust at the rate of 35 lbs. per acre, equal to 1-3/4 oz. per 50 feet of row.

**BEANS—**For the control of Mexican bean beetles, leafhoppers, spider mites, Japanese beetles, aphids, cucumber beetles. Do not harvest within one day of application: Apply 30 to 35 lbs: per acre.

**COLE CROPS & LEAFY VEGETABLES—**Broccoli, Brussels sprouts, cabbage, collards, kale, mustard greens, parsley, turnips. For the control of aphids. Do not harvest within 7 days of application. Apply 30 to 35 pounds per acre.

**CUCURBITS—**Cucumber, melons. Do not apply within one day of harvest. For the control of aphids, pickle worms, leafhoppers, apply 32 to 35 lbs per acre.

**LETTUCE AND SPINACH—**For the control of aphids: Do not apply to head lettuce within 7 days of harvest or leaf lettuce within 14 days of harvest.

**PEAS—**For the control of pea aphids, pea weevils: Do not apply within 7 days of harvest. Apply 30 to 35 lbs per acre.

**POTATOES—**For the control of aphids, leafhoppers, mealy bugs. Apply 30 to 35 lbs per acre.

**TOMATOES—**For the control of aphids. Do not apply within one day of harvest. Apply 35 pounds per acre.

**POULTRY HOUSES—**For the control of Northern Fowl Mites, Chicken Red Mite and Poultry Lice, apply 3/4 pound per 50 to 60 square feet of litter, floor space, nests, roosts and adjacent areas. Repeat as necessary.

**CATTLE—**For the control of lice on beef cattle make a thorough application to the animals. Repeat application within 10 days and thereafter as needed. Avoid contamination of feed, food containers and watering troughs. Do not treat lactating dairy animals. Do not apply later than 5 hours before milking or during milking. Do not treat calves under one month of age.

**SWINE—**For the control of lice on swine, make a thorough application of the animals. Pens should be dusted: Repeat application in 10 days and thereafter as needed: Do not treat animals under 1 month of age.

## CAUTION

### KEEP OUT OF THE REACH OF CHILDREN

**AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. HARMFUL IF SWALLOWED.**

**NOTE TO PHYSICIANS:** Malathion may cause cholinesterase inhibition. Atropine is antidotal.

#### — ENVIRONMENTAL CAUTION —

This product is toxic to fish. Keep out of lakes, streams, or ponds: Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops: Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not contaminate feed, food containers and water troughs.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

#### NOTICE

Steven. Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Stevens Industries, Inc. and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.

MANUFACTURED BY

**STEVENS INDUSTRIES, INC., DAWSON, GEORGIA 31742**

EPA Est No 2459-GA-1

EPA Reg. No. 2459-181