

BEST DOCUMENT AVAILABLE
CAUTION

KEEP OUT OF THE REACH OF CHILDREN

DIRECTIONS

For The Control of:
BITES: LEAFHOPPERS TOMATO RUSSET MITES
LACE BUGS: APHIDS, THRIPS MEXICAN BEAN BEETLES
SPIDER MITES: CABBAGE LOOPERS IMPORTED CABBAGE WORMS

Attaching:
BEANS: TURNIPS MELONS PEAS
CABBAGE: CUCUMBERS EGGPLANT PEPPERS
BROCCOLI: SQUASH ONIONS TOMATO

To control these insects, dust at the rate of 25 to 30 pounds per acre or 2 ounces to each 50 ft. of row in the garden. Repeat applications at 5 to 7 day intervals, as necessary for control. Do not apply to plants unless they are dry or injury may result. Do not use on beans, cucumbers, melons, or squash within 1 day of harvest; on broccoli, eggplant, onions, peas, peppers, tomatoes, turnips or potatoes within 3 days of harvest; on cabbage within 7 days of harvest.

POULTRY

Chickens, ducks, turkey and geese: To control Northern fowl mite, lice, and as a supplement to premise treatment for chicken red mite. Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 6 weeks or when necessary.

Premise Treatments: To control Northern fowl mite, chicken red mite and poultry lice apply at the rate of 1 lb. to 50-60 square feet of litter and floor space, to nests, roosts and adjacent areas. Repeat application as necessary. AVOID CONTAMINATION OF FEED AND FEEDING TROUGHS AND WATER FOUNTAINS.

DOGS AND CATS

Fleas: Apply just enough dust to treat animal thoroughly. Apply to the back and work into the coat with the hand. Wash hands thoroughly after using. Repeat applications as necessary for control but not more often than once weekly.

ORNAMENTALS

Aphids, red spider and bagworm: Dust infested plants thoroughly. Repeat applications as necessary for control. NOTE: Injury may result on certain ferns including Boston, Maidenhair and Pteris and on some species of Crassula.

CAUTION

May be harmful if swallowed. Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Do not contaminate ornamental fish ponds. Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

NOTE TO PHYSICIANS

Malathion, upon repeated, prolonged, or careless use, may cause Cholinesterase inhibition. Atropine is antidotal.

U.S.D.A REG. NO. 242-122

14

14A



CONTENTS

NET WT.

1 POUND

5%

NOV-4-1968

2459 183

**MALATHION
DUST**

DANGER

KEEP OUT OF THE REACH OF CHILDREN
SEE OTHER PANEL FOR ADDITIONAL WARNINGS

THIS PACKAGE CONTAINS

ACTIVE INGREDIENTS:	5.0%	Malathion (O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate).
INERT INGREDIENTS	95.0%	
Total	100.0%	

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to Stevens Industries, Inc., and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.

Manufactured by

STEVENS INDUSTRIES, INC., Dawson, Georgia

21742

31A

31

BEST DOCUMENT AVAILABLE