

ACCEPTED
APR 24 1968
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PESTS REGISTERED
 UNDER NO. 257-150000
 TO ATTACHED LABEL



BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

MALATHION EMULSIFIABLE SPRAY
ONE GALLON CONTAINS FIVE POUNDS OF MALATHION
 CONTENTS BY WEIGHT

ACTIVE INGREDIENTS:	
Malathion*	57%
Xylene	32%
INERT INGREDIENTS:	
	11%
TOTAL	100%

CAUTION
KEEP OUT OF REACH OF CHILDREN

CAUTION
KEEP OUT OF REACH OF CHILDREN

* O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

RECOMMENDATIONS

COTTON: For the control of Boll Weevil, Cotton Aphid, Cotton Leaf Worm, Desert Spider Mite, use 1 2/3 - 3 1/3 pints per acre with sufficient water to cover one acre. Consult State Agricultural Experiment Station, or State Extension Service Specialists for time and number of applications.

VEGETABLES: For the control of Aphids, Spider Mites, Pickleworms and Cucumber beetle attacking CUCUMBERS, SQUASH and MELONS; and the control of Mexican Bean Beetles, Spider Mites, Aphids attacking BEANS and PEAS; and the control of Aphids, Cabbage Worms, Cabbage Looper, Flea Beetles, Spider Mites attacking CABBAGE, TURNIPS, ONIONS, SPINACH, MUSTARD, LETTUCE, POTATOES and TOMATOES. Use 1 1/2-2 pints per acre in sufficient water for good coverage. Repeat application at 5 to 7 day intervals, as necessary for control. Do not use on cabbage, spinach, mustard within 7 days of harvest. Do not use on other vegetables named within 3 days of harvest. Do not apply to leaf lettuce within 14 days of harvest.

FRUITS: For control of Aphids, Red Spider Mites, Mealy Bugs, Woolly Aphids, White Fly, Thrips, Tarnished Plant Bug, Pear Psyllid, Scale Crawlers, Plum Curculia attacking Apples, Pears, Grapes, Peaches and Plums.

Use 1 1/2-2 teaspoonfuls per one gallon of water. Repeat applications as necessary. Malathion may cause injury to McIntosh and Courtland varieties of apples and Ribier grapes. Do not apply within 3 days of harvest. Do not use on peaches within 7 days of harvest.

ORNAMENTALS: For the control of Aphids, Red Spider Mites, Woolly Aphids, White Fly, Thrips, Roseleaf Hopper attacking roses, camellias, azaleas and evergreens. Use 1 1/2 to 2 teaspoonfuls per one gallon of water. Do not use on Boston, Maidenhair and Petrus Ferns as well as some species of Crassula.

FLIES AND MOSQUITOES: Spray around windows, door sills of houses, crevices and woodwork in outhouses and barns. Mix one part of Malathion to 28 parts of water. Use one gallon of diluted spray to cover 1000 sq. ft. Avoid contamination of milk, milking equipment and water.

CAUTION

Harmful if swallowed. Flammable. Avoid prolonged breathing of spray mist. Avoid repeated or prolonged contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs. Keep container closed and away from heat and open flame. Highly toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.

NOTE TO PHYSICIANS: Malathion, upon repeated, prolonged, careless use may cause Cholinesterase Inhibitor.. ATROPINE IS ANTIDOTAL.

NOTICE

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Stevens Industries, Inc., and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.

USDA Reg. No. 2450-189

Manufactured By
STEVENS INDUSTRIES, INC.
 DAWSON, GEORGIA 31708