

NET WEIGHT 50 LBS.



**master**®

B R A N D

**2% ETHYL PARATHION DUST**

THIS PACKAGE CONTAINS

ACTIVE INGREDIENTS:

Parathion (O,O-Diethyl O-p-nitrophenyl thiophosphate)

2.00%

INERT INGREDIENTS:

98.00%

TOTAL

100.00%

DIRECTIONS

**PEAS, BEANS, CABBAGE, COLLARDS, LETTUCE, MUSTARD, TURNIPS:** For aphid control, apply 12.5 Tbs. Master Brand 2% Ethyl Parathion per acre. Inspect vegetables frequently for insects and insect damage so that the insecticides can be applied before damage becomes severe. Applications of insecticides should be made every two weeks or oftener if needed. It is usually necessary to apply insecticides more often during periods of rainy weather and heavy infestations. Be sure to dust thoroughly at lay-by time.

CAUTIONS

Peas	Do not apply 10 days prior to harvest	Head Lettuce	Do not apply 7 days prior to harvest
Beans	Do not apply 15 days prior to harvest	Bibb or Leaf Lettuce	Do not apply 21 days prior to harvest
Cabbage	Do not apply 10 days prior to harvest	Mustard	Do not apply 10 days prior to harvest
Collards	Do not apply 10 days prior to harvest	Turnips	Do not apply 10 days prior to harvest

**WARNING**

**KEEP OUT OF REACH OF CHILDREN!**

Hazardous if swallowed, inhaled, or rapidly absorbed through skin. Avoid inhalation and skin contact. Wear protective clothing, natural rubber gloves and goggles. Wear an effective toxic respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Wash thoroughly with soap and warm water after exposure. Don't keep poisons where food is handled or stored. Avoid contamination of feed and foodstuffs. Keep all persons out of the operating area or vicinity where there may be danger of drift. Vacated areas should not be entered until the drifting insecticides and volatile residues have dissipated.

**CAUTION:** Workers entering field within 24 hours of application should wear protective clothing.

Do not apply when weather conditions favor drift of dust from crop area treated.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

applied before damage becomes severe. Applications of insecticides should be made every two weeks or oftener if needed. It is usually necessary to apply insecticides more often during periods of rainy weather and heavy infestations. Be sure to dust thoroughly at lay-by time.

### CAUTIONS

Peas	Do not apply 10 days prior to harvest	Head lettuce	Do not apply 7 days prior to harvest
Beans	Do not apply 15 days prior to harvest	Bibb or leaf lettuce	Do not apply 21 days prior to harvest
Cabbage	Do not apply 10 days prior to harvest	Mustard	Do not apply 10 days prior to harvest
Collards	Do not apply 10 days prior to harvest	Turnips	Do not apply 10 days prior to harvest

## WARNING

### KEEP OUT OF REACH OF CHILDREN!

Hazardous if swallowed, inhaled, or rapidly absorbed through skin. Avoid inhalation and skin contact. Wear protective clothing, natural rubber gloves and goggles. Wear an effective toxic respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Wash thoroughly with soap and warm water after exposure. Don't keep poisons where food is handled or stored. Avoid contamination of feed and foodstuffs. Keep all persons out of the operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until the drifting insecticides and volatile residues have dissipated.

**CAUTION:** Workers entering field within 24 hours of application should wear protective clothing.

Do not apply when weather conditions favor drift or dust from crop area treated.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.



**INTERNAL:** Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep warm.

**EXTERNAL:** If the skin is contaminated, wash thoroughly with soap and water.

### ANTIDOTE FOR PARATHION

**CALL A PHYSICIAN AT ONCE!** (Note: For physician only Therapeutic doses of atropine appear to be effective as an antidote.)

**BAGS:** Completely empty bag and dispose of waste pesticide by burying. Bury bag and waste pesticide at least 18 inches deep in an isolated location away from water supplies.

### NOTICE

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damage such as loss of profits.

MANUFACTURED BY

**STEVENS INDUSTRIES, INC.**

**DAWSON, GEORGIA 31742**

U.S.D.A. Reg. No. 2458-200