

NET WEIGHT 50 LBS



Master

BRAND

6% DITHANE M-45 DUST

A FUNGICIDE DUST

ACTIVE INGREDIENTS

A coordination product of zinc and manganese ethylene bisdithiocarbamate
In which the ingredients are:

| | |
|-----------------------------|-----|
| Manganese | 96 |
| Zinc | 12 |
| Ethylene bisdithiocarbamate | 172 |

INERT INGREDIENTS

TOTAL

100.0

CAUTION

KEEP OUT OF REACH OF CHILDREN!

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes, wash copiously with water and get medical attention. Do not reuse containers. Destroy by burying with waste. Contamination of fish-bearing waters with this product may harm fish.

IMPORTANT: Store in cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of this product. Keep container closed when not in use.

DIRECTIONS FOR USE

Peanuts: Cercospora leaf spot: Begin applications when disease first appears and repeat at 10-14 day intervals throughout the season or as required for adequate disease control. Apply 20-35 lbs. acre as required to provide good coverage of plants. If peanut hay is to be used as livestock feed, Dithane M-45 dust should be applied at a dosage of 20-25 lbs./acre and should not be applied within 14 days of harvest.

Potatoes: For early and late blight, apply every 7-10 days throughout the season. Begin application when plants are 4 to 6 inches high, applying 20 pounds per acre and increasing to 25 pounds as the vines increase in size.

Tomatoes: Late blight, early blight, Septoria and gray leaf spots, gray leaf mold and anthracnose. Begin application when plants are 4 to 6 inches high, applying 20 pounds per acre and increasing to 25 pounds as the vines increase in size. Make applications at 7 day intervals.

Cucumbers, Summer Squash, Cantaloupes, Honeydew Melons, Muskmelons and Watermelons: Downy mildew, anthracnose, scab, alternaria leaf spot, Cercospora leaf spot and gummy stem blight. Begin applications at the two leaf stage and repeat at 5 to 7 day intervals, applying 20 pounds per acre and increasing to 30 pounds per acre as the vines increase in size.

MASTER BRAND 6% DITHANE M-45 DUST will suppress the diseases listed by the respective crops, however, to achieve the desired results, a regular schedule must be followed and applications must be applied under conditions favorable for the use of dust.

CAUTION

KEEP OUT OF REACH OF CHILDREN!

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray. In case of skin contact wash with water; for eyes, wash copiously with water and get medical attention. Do not reuse containers. Destroy by burying with waste. Contamination of fish-bearing waters with this product may harm fish.

IMPORTANT! Store in cool, dry place. Do not allow to become wet or overheated. This may bring on chemical changes which will impair the fungicidal effectiveness of this product. Keep container closed when not in use.

DIRECTIONS FOR USE

Peanuts Cercospora leaf spot: Begin applications when disease first appears and repeat at 10-14 day intervals throughout the season or as required for adequate disease control. Apply 20-35 lbs./acre as required to provide good coverage of plants. If peanut hay is to be used as livestock feed, Dithane M-45 dust should be applied at a dosage of 20-25 lbs./acre and should not be applied within 14 days of harvest.

Potatoes: For early and late blight, apply every 7-10 days throughout the season. Begin application when plants are 4 to 6 inches high, applying 20 pounds per acre and increasing to 25 pounds as the vines increase in size.

Tomatoes: Late blight, early blight, Septoria and gray leaf spots, gray leaf mold and anthracnose. Begin application when plants are 4 to 6 inches high, applying 20 pounds per acre and increasing to 25 pounds as the vines increase in size. Make applications at 7 day intervals.

Cucumbers, Summer Squash, Cantaloupes, Honeydew Melons, Muskmelons and Watermelons: Downy mildew, anthracnose, scab, alternaria leaf spot, Cercospora leaf spot and gummy stem blight. Begin applications at the two leaf stage and repeat at 5 to 7 day intervals, applying 20 pounds per acre and increasing to 30 pounds per acre as the vines increase in size.

MASTER BRAND 6% DITHANE M-45 DUST will suppress the diseases listed by the respective crops, however, to achieve the desired results, a regular schedule must be followed and applications must be applied under conditions favorable for the use of dust.

DITHANE is a trademark Reg. in U. S. Pat. Off. and in principal foreign countries by Rhom and Haas Company.

NOTICE

Stevens Industries, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Stevens Industries, Inc., and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damage such as loss of profits.

MANUFACTURED BY

STEVENS INDUSTRIES, INC.

DAWSON, GEORGIA 31742

U.S.D.A. Reg. No. 2459-208