



likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

- VEGETABLES -

Use for control of surface-feeding weevils, cutworms, and other insects, army worms and crickets on the crops listed on this label. Apply before or just after plowing fields with a history of surface feeding; cutworms, army worms and crickets on these fields suspected of infestation.

PEAS, BEANS: Use 15 pounds per acre. Do not apply within 21 days of harvest.

CORN, SOYBEAN, PEA, BEAN, CUCURBIT: Use 30 pounds per acre. Apply only 2 applications per season. Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy cattle.

CUCURBIT, LETTUCE: Use 25 pounds per acre. Do not apply within 21 days of harvest. Do not apply on lettuce after heads begin to form.

POTATO, SWEET POTATO, CABBAGE, CARNIFLOWER: Use 25 pounds per acre. Do not apply within 21 days of harvest.

CELERY: Use 30 pounds per acre. Do not apply within 30 days of harvest. Do not spray more than twice per season.

SPINACH: (dry seed crop) Use 25 pounds per acre. Do not apply within 14 days of harvest. Do not spray more than twice per season. Do not feed to meat or dairy cattle.

- FRUIT CROPS -

Use for control of surface-feeding weevils, cutworms, and other insects on the crops listed on this label. Apply before or just after plowing fields with a history of surface feeding; cutworms, army worms and crickets on these fields suspected of infestation.

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.