


# HALCO

## GRANULAR TURF FUNGICIDE



For the prevention and control of COPPER SPOT, DOLLAR SPOT, LEAF SPOT (Helminthosporium-Curvularia), BROWN PATCH and SNOW MOLD on Lawns containing Kentucky Blue including Merion and other strains, the various Fescues, Bents and most other grasses.

### SPECIAL INSTRUCTIONS:

Do not water or mow lawn area for 24 hours after application. Application will be more effective if made in the early morning when dew is on the grass. Repeat treatments are necessary until fungus diseases disappear. Apply contents of this bag to 10,000 square feet for prevention program and at double rate (5000 square feet) if fungus diseases have already infested lawn area.

### SPRING PROGRAM:

Application should be made immediately at a double rate (contents of this bag applied to 5000 square feet) if no disease treatments have been given and Dollar or Leaf Spot have invaded Lawn area. For a prevention program against these diseases apply HALCO GRANULAR TURF FUNGICIDE soon after the second mowing in the Spring and repeat every 7 to 10 days for a minimum of five times. This program will prevent Fungus activity of Dollar Spot and Leaf Spot which may kill good grasses.

### SUMMER APPLICATION:

Apply immediately at double rate if Brown Patch or Curvularia Disease is active in lawn area. Prevention of these diseases may be obtained by an application of HALCO GRANULAR TURF FUNGICIDE every 7 to 10 days during the hot humid months.

### EARLY FALL APPLICATION:

Prevent infestation of Dollar Spot and Copper Spot with treatments at 7 to 10 day intervals during the period of cooler nights.

### SNOW MOLD:

Apply at double rate in the late Fall, before the first snowfall and repeat again at double rate in January.

## CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contact with eyes, skin or clothing. In case of contact wash exposed areas thoroughly with water and avoid irritation. Avoid breathing dust since inhalation may prove harmful.

**INTERNAL:** If swallowed give raw white of egg. Induce vomiting with mixture of salt, soap or mustard in warm water. Call physician at once.

Consumption of alcoholic beverages increases the possibility of harm.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish and wildlife, keep out of lakes, streams or ponds. Do not contaminate water by cleaning out equipment, or disposal of wastes.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and or handling is contrary to label instructions.

### Spreader Settings for Proper Application —

Normal Rate of 20 Pounds per 10,000 Square Feet

Scotts	5 1/2	Buch #2 - 20"	3/4
Lawn Beauty	5	Cyclone	3.4
Jackson - 20"	30	Others	3 1/2 o. an inch opening

Where Double Rate application is wanted, go over area twice at above settings.

APPLY with SPREADER      COVERS 10,000 SQUARE FEET

**ACCEPTED**

**MAR 13 1972**

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT AND REGULATIONS FOR ECONOMIC PESTICIDE MANAGEMENT ORDER NO. 2772-2

For the prevention and control of **COPPER SPOT, DOLLAR SPOT, LEAF SPOT** (Helminthosporium-Curvularia), **BROWN PATCH** and **SNOW MOLD** on Lawns containing Kentucky Blue including Merion and other strains, the various Fescues, Bents and most other grasses.

**SPECIAL INSTRUCTIONS:**

Do not water or mow lawn area for 24 hours after application. Application will be more effective if made in the early morning when dew is on the grass. Repeat treatments are necessary until fungus diseases disappear. Apply contents of this bag to 10,000 square feet for prevention program and at double rate (5000 square feet) if fungus diseases have already infested lawn area.

**SPRING PROGRAM:**

Application should be made immediately at a double rate (contents of this bag applied to 5000 square feet) if no disease treatments have been given and Dollar or Leaf Spot have invaded Lawn area. For a prevention program against these diseases apply HALCO GRANULAR TURF FUNGICIDE soon after the second mowing in the Spring and repeat every 7 to 10 days for a minimum of five times. This program will prevent Fungus activity of Dollar Spot and Leaf Spot which may kill good grasses.

**SUMMER APPLICATION:**

Apply immediately at double rate if Brown Patch or Curvularia Disease is active in lawn area. Prevention of these diseases may be obtained by an application of HALCO GRANULAR TURF FUNGICIDE every 7 to 10 days during the hot humid months.

**EARLY FALL APPLICATION:**

Prevent infestation of Dollar Spot and Copper Spot with treatments at 7 to 10 day intervals during the period of cooler nights.

**SNOW MOLD:**

Apply at double rate in the late Fall, before the first snowfall and repeat again at double rate in January.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

Avoid contact with eyes, skin or clothing. In case of contact wash exposed areas thoroughly with water and avoid irritation. Avoid breathing dust since inhalation may prove harmful.

**INTERNAL:** If swallowed give raw white of egg. Induce vomiting with mixture of salt, soap or mustard in warm water. Call physician at once.

Consumption of alcoholic beverages increases the possibility of harm.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish and wildlife, keep out of lakes, streams or ponds. Do not contaminate water by cleaning out equipment, or disposal of wastes.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**Spreader Settings for Proper Application —**

Normal Rate of 20 Pounds per 10,000 Square Feet

Scotts	5 1/2	Buch #2 - 20"	3/4
Lawn Beauty	5	Cyclone	3.4
Jackson - 20"	30	Others	3/16 of an inch opening

Where Double Rate application is wanted, go over area twice at above settings.

**APPLY with SPREADER      COVERS 10,000 SQUARE FEET**

**ACTIVE INGREDIENTS**

Thiram (Tetramethylthiuram Disulfide) ..... 5.00%

\*Cadmium Chloride ..... .75%

**INERT INGREDIENTS** ..... 94.25%

\*Total Cadmium ..... .43%

**NET WEIGHT 20 POUNDS**

EPA  
Reg. No.  
2474-47

MANUFACTURED FOR  
**HALCO DIVISION OF WAGNER SEED COMPANY, INC.**  
81 SECATOGUE A.V., FARMINGDALE, N. Y. 11735