

3491-227

HOLIDAY CATTLE FLY SPRAY

DIRECTIONS FOR USE

To develop a residual repellency to stable flies, horn flies, mosquitos and gnats on dairy cows, apply 1 to 2 ounces of HOLIDAY CATTLE FLY SPRAY to each animal for a three day period. The spray should be applied with a good sprayer so that the entire animal's body has been uniformly covered. Special care should be used in spraying the legs, belly and shoulders. Avoid areas around the nose, eyes and mouth. Avoid wetting the hide. Spray in the same manner again after two days have elapsed and every second day thereafter or when insects again disturb the animals. Be sure the teats and udder are washed with warm water prior to milking. Calves should be sprayed very lightly once a day instead of attempting to build up a residual repellency by spraying more heavily.

For control of lice on dairy and beef animals, spray the animals with 2 ounces of HOLIDAY CATTLE FLY SPRAY. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

To kill flies, mosquitoes and gnats in the barn prior to milking, close all doors and windows and direct the spray toward the ceiling. The barn should remain closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary.

To protect dairy and beef animals from attack by fleas, spray bedding and walls of pen particularly cracks and crevices. Repeat treatment frequently until adequate control is obtained.

In a complete fly control program these sprays should be used in combination with residual insecticides and sanitary measures.

CAUTION:

Harmful if swallowed. Avoid contamination of milk and milk handling equipment. Avoid overspraying which will result in wetting the animal's hide. Do not spray on or near open flame. Store in cool place. Keep container tightly closed. Do not refill this container.

ANTIDOTE — INTERNAL: Use an emetic of mustard to induce vomiting. Call your physician. If splashed in eye, flush thoroughly with water.

READY-TO-USE

HOLIDAY

CATTLE FLY SPRAY

MADE WITH MGK CONCENTRATE 933 AND MGK REPELLENT 326

Kills and repels horn flies, stable flies, mosquitoes and gnats. When properly applied, the repellent effect against above insects will last two days.

ACTIVE INGREDIENTS:

Pyrethrins	0.3%
*Piperonyl butoxide, technical	0.70
Nonyl bis(2-hydroxyethyl)benzene oxide	1.15
*Dimethyl heptadecylbenzoate	2.00
Petroleum distillate	99.580

Pyrethrins, Piperonyl butoxide, Nonyl bis(2-hydroxyethyl)benzene oxide, Dimethyl heptadecylbenzoate, Petroleum distillate. © 1968 Koos Inc., Kenosha, Wisconsin 53140. Manufactured by Koos Inc., Kenosha, Wisconsin 53140. MGK Concentrate 933 and MGK Repellent 326. U.S.D.A. Reg. No. 2491-221.

ONE GALLON

CAUTION: "Keep out of the reach of children."

(SEE OTHER WARNINGS ON SIDE PANELS)

KOOS INC., KENOSHA, WISCONSIN U.S.A. 53140

U.S.D.A. REG. NO. 2491-221

HOLIDAY CATTLE FLY SPRAY

DIRECTIONS FOR USE

To develop a residual repellency to stable flies, horn flies, mosquitos and gnats on dairy cows, apply 1 to 2 ounces of HOLIDAY CATTLE FLY SPRAY to each animal for a three day period. The spray should be applied with a good sprayer so that the entire animal's body has been uniformly covered. Special care should be used in spraying the legs, belly and shoulder. Avoid areas around the nose, eyes and mouth. Avoid wetting the hide. Spray in the same manner again after two days have elapsed and every second day thereafter or when insects again disturb the animals. Be sure the teats and udder are washed with warm water prior to milking. Calves should be sprayed very lightly once a day instead of attempting to build up a residual repellency by spraying more heavily.

For control of lice on dairy and beef animals, spray the animals with 2 ounces of HOLIDAY CATTLE FLY SPRAY. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

To kill flies, mosquitoes and gnats in the barn prior to milking, close all doors and windows and direct the spray toward the ceiling. The barn should remain closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary. To protect dairy and beef animals from attack by fleas, spray bedding and walls of pen particularly cracks and crevices. Repeat treatment frequently until adequate control is obtained. In a complete fly control program these sprays should be used in combination with residual insecticides and sanitary measures.

CAUTION:

Harmful if swallowed. Avoid contamination of milk and milk handling equipment. Avoid overspraying which will result in wetting the animal's hide. Do not spray on or near open flame. Store in cool place. Keep container tightly closed. Do not refill this container.

ANTIDOTE — INTERNAL: Use an emetic of mustard to induce vomiting. Call your physician. If splashed in eye, flush thoroughly with water.

