

HOLIDAY

SEVIN*—5%

Home and Garden Dust

For Control of Certain Insects on Vegetables,
Ornamentals and Pets

Active Ingredient:	by Wt.
Carbaryl (1-naphthyl N-methylcarbamate).....	5%
Inert Ingredients	95%
Total	100%

U.S. Pat. No. 2,903,478; 3,009,855

Read entire label and use in strict accordance with label directions and cautions.

General Directions: Apply dust when insects or damage appears on plants. Make applications in the morning or evening when it is cool and the air is still. Treat upper and lower surface of foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results apply with a dust gun. Do not apply in excess.

NET WT. 1 LB.

1900 Third Street / Kenosha, Wisconsin 53140

READY-TO-USE

VEGETABLE INSECT CONTROL

Asparagus—To control asparagus beetle and Apache cicada. Do not treat within one day of harvest. Do not repeat applications at intervals less than 3 days apart.

Beans (green beans, lima beans, snap beans, cowpeas, black-eyed peas)—To control Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworms, corn earworm, Lima bean pod borer, stink bugs and tarnished plant bug.

Sweet Corn—To control corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetles, sap beetles and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.—For insects attacking silks and ears, apply at 2-3 day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

Cucumber, Melons, Pumpkin, Squash—To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. SEVIN injures watermelons in Florida.

Tomato, Eggplant, Pepper, Peas—To control Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

Carrots, Horseradish, Head Lettuce, Parsnips, Radishes, Rutabagas, Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi—To control flea beetles, harlequin bug, leafhoppers, armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs and stink bugs. To control small cabbage loopers, use on a 5 to 7-day schedule. Do not treat within 3 days of harvest.

Chinese Cabbage, Collards, Dandelion, Endive (escarole), Garden Beets, Kale, Mustard Greens, Parsley, Salsify, Spinach, Swiss Chard and Turnips—To control flea beetles, harlequin bug, leafhoppers, armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs and stink bugs. To control small cabbage loopers, use on a 5 to 7 day schedule. Do not treat within 14 days of harvest.

ORNAMENTAL INSECT CONTROL

This product may be safely used on a wide range of ornamental plants, flowers, shrubs, and small trees such as rose, carnation, gladiolus, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not apply to Boston Ivy. Apply to the upper and lower leaf surface of all foliage for complete coverage to control such pests as: apple aphid, birch leaf miner, boxelder bug, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetle, leafhoppers, leaf roller, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles. Note: Sevin does not control spider mites. For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop or neighboring crops. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

PET USE

On dogs and cats: For ticks or fleas dust liberally over the entire animal, avoiding eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Do not treat kittens or puppies less than 4 weeks of age. In and around dog and cat quarters: For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

Do not breathe dust if swallowed or inhaled. Avoid prolonged or repeated breathing of dust. Do not take internally. Avoid contact with skin—wash hands and face after use. Do not contaminate feed or foodstuffs.

*Registered Trade Mark of Union Carbide Corporation for the active ingredient (1-naphthyl N-methylcarbamate)

KOOS, INC.

Kenosha, Wisconsin 53140

ACCEPTED

July 9, 1970

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 2491-234 SUBJECT
TO ATTACHED COMMENTS

U.S.D.A. REG. NO. 2491-234

Product of U.S.A.