

# HOLIDAY TOMATO VEGETABLE DUST

Controls many insects and diseases on  
Tomatoes, Melons, Squash, Potatoes,  
and Certain other Vegetables  
and Small Fruits

Active Ingredients:	By Wt.
Captan*	5.00%
Methoxychlor Technical**	5.00%
Rotenone from Cube	0.75%
Other Resins from Cube	0.75%
Inert Ingredients:	88.50%
Total	100.00%

\*N-[(trichloromethyl) thio]-4-cyclohexene-2,2-dicarboximide.  
\*\*Equivalent to 4.4% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% other isomers and related compounds.  
U. S. Pat. No.'s 2,553,770-2,553,771

NET WT. 10 OZ.

# HOLIDAY TOMATO - VEGETABLE DUST

Read complete label and use in strict accordance with directions, warnings and cautions.

For use on tomatoes and other vegetables and certain small fruits.

**Directions:** Apply dust in the morning or evening when it is cool and the air is still. Treat upper and lower surface of foliage for complete coverage. Repeat as necessary or at 5-10 day intervals.

This product may be used as a spray by mixing 3 to 4 tablespoons of this material in a little water and forming a thick paste. Then add to 1 gallon water and keep mixed while spraying. Do not store spray but make new solution each time application is desired.

Apply when insects first appear.

Controls the following insects on:

Tomatoes—aphids, thrips, leafhoppers, flea beetles, blister beetles, tomato fruit worms.

Beans—aphids, thrips, leafhoppers.

Squash—aphids, leafhoppers, squash bug (nymphs) blister beetles, squash vine borers.

Eggplant—aphids, flea beetles, blister beetles.

Watermelon—aphids, leafhoppers, thrips, blister beetles, squash bug (nymphs).

Cantaloupes—aphids, thrips, leafhoppers, blister beetles, squash bug (nymphs).

Potatoes—aphids, leafhoppers, flea beetles, blister beetles, Colorado potato beetle.

Peppers—thrips, flea beetles.

Strawberries—aphids, leafhoppers, spittle bugs.

Raspberries—aphids, raspberry fruitworms.

Also controls—melon and pickle worms, cucumber beetles, Japanese beetles and Mexican bean beetles.

For control of diseases: When leaves begin to unfurl start dusting. Repeat at 1-2 week intervals. Controls Tomato and Potato Blight, Anthracnose, Tomato Septoria Leaf Spot, Alternaria, Cercospora of Peppers, Phomopsis, Botrytis rot of strawberries, Fruit rot of raspberries, and Spur blight.

May be used on beans and potatoes up to within 3 days of harvest but do not use on other crops within 7 days of harvest.

Toxic to fish.

Do not contaminate ponds, lake, or streams or water used for domestic purposes with this product.

**CAUTION:** Do not breathe dust or spray. Avoid contact with skin. Wash hands after use. Do not contaminate feed and foodstuffs. Wash fruits and vegetables before using.



4500 Thirteenth Court / Kenosha, Wisconsin 53140

ACCEPTED  
Mar 16, 1970

U.S.D.A. REG. NO. 2491-236

2491-236



# HOLIDAY

## TOMATO VEGETABLE DUST

Controls many insects and diseases on  
Tomatoes, Melons, Squash, Potatoes,  
and Certain other Vegetables  
and Small Fruits

Active Ingredients:  
Captan\*  
Methoxychlor Technical\*\*

By Wt.  
5.00%  
5.00%

# HOLIDAY

## TOMATO - VEGETABLE DUST

Read complete label and use in strict accordance with directions, warnings and cautions.

For use on tomatoes and other vegetables and certain small fruits.

**Directions:** Apply dust in the morning or evening when it is cool and the air is still. Treat upper and lower surface of foliage for complete coverage. Repeat as necessary or at 5-10 day intervals.

This product may be used as a spray by mixing 3 to 4 tablespoons of this material in a little water and forming a thin paste. Then add to 1 gallon water and keep mixed while spraying. Do not store spray but make new solution each time application is desired.

Apply when insects first appear. —

Controls the following insects on:

Tomatoes—aphids, thrips, leafhoppers, flea beetles, blister beetles, tomato fruit worms.

Beans—aphids, thrips, leafhoppers.

Squash—aphids, leafhoppers, squash bug (nymphs) blister beetles, squash vine borers.

Eggplant—aphids, flea beetles, blister beetles.

Watermelon—aphids, leafhoppers, thrips, blister beetles, squash bug (nymphs).

Cantaloupes—aphids, thrips, leafhoppers, blister beetles, squash bug (nymphs).

Potatoes—aphids, leafhoppers, flea beetles, blister beetles, Colorado potato beetle.

Peppers—thrips, flea beetles.

Strawberries—aphids, leafhoppers, spittle bugs.

Raspberries—aphids, raspberry fruitworms.

Also controls—melon and pickle worms, cucumber beetles, Japanese beetles and Mexican bean beetles.

For control of diseases: When leaves begin to unfurl start dusting. Repeat at 1-2 week intervals. Controls Tomato and Potato Blight, Anthracnose, Tomato Septoria Leaf Spot, Alternaria, Cercospora of Peppers, Phomopsis, Botrytis rot

# VEGETABLE DUST

Controls many insects and diseases on  
Tomatoes, Melons, Squash, Potatoes,  
and Certain other Vegetables  
and Small Fruits

Active Ingredients	By Wt.
Captan*	5.00%
Methoxychlor Technical**	5.00%
Rotone from Cuba	0.75%
Other Resins from Cuba	0.75%
Inert Ingredients:	88.50%
Total	100.00%

\*N-(1-trichloromethyl-4-piperidyl)phthalic di-  
carboximide

\*\*Equivalent to 4.4% 2,2-bis(4-methoxyphenoxy)-  
1,1,1-trichloroethane and 0.6% other  
isomers and related compounds.  
U.S. Pat. Nos. 2,553,770; 2,553,771

NET WT. 10 OZ.

Squash—aphids, leafhoppers, squash bug, nymphs, blister beetles, squash  
vine borers

Eggplants—aphids, flea beetles, blister beetles

Watermelons—aphids, leafhoppers, thrips, blister beetles, squash bug, nymphs

Cantaloupes—aphids, thrips, leafhoppers, blister beetles, squash bug, nymphs

Potatoes—aphids, leafhoppers, flea beetles, blister beetles, Colorado potato  
beetle

Peppers—thrips, flea beetles

Strawberries—aphids, leafhoppers, spittle bugs

Raspberries—aphids, raspberry fruit twerms

Also controls melon and pink worms, cucumber beetle, Japanese beetles  
and Mexican bean beetle

For control of diseases: When leaves begin to unfurl start dusting. Repeat at  
1-2 week intervals. Controls Tomato and Potato Blight, Anthracnose, Tomato  
Septoria Leaf Spot, Alternaria, Cercospora of Peppers, Phomopsis, Botrytis rot  
of strawberries, fruit rot of raspberries, and Spur blight.

May be used on leeks and potatoes up to within 7 days of harvest, but do not  
use on other crops within 7 days of harvest.

Toxic to fish

Do not contaminate water, streams, or water used for domestic pur-  
poses with this product.

**CAUTION:** Do not breathe dust or spray. Avoid contact with skin.  
Wash hands after use. Do not contaminate feed and foodstuffs.  
Wash fruit and vegetables before using.

4500 Thirteenth Court / Kenosha, Wisconsin 53140

U.S.D.A. REG. NO. 2491-236