

SUGAR BAIT FLY KILLER

For the control of—Flies around Dairy Barns,
Horse Barns—Stock Barns—Pig Pens—
Kennels—Outdoor Living Areas—
Milk Sheds

Active Ingredients	By Wt.
2,2-Dimethyl-3-cyanoethyl dimethyl phosphorothioate	0.92%*
Related compounds	0.08%*
Inert ingredients	99.00%
Total	100.00%

*Registered under FIFRA of the U.S. Department of Agriculture
© Trademark of J. Kenosha Chemical Company

Net Contents:
1 Pound

Directions for Use & Application

Dairy Barns, Milk Sheds, Stables, Livestock Barns, Loafing Shed, Pig Pens, & Outdoor Areas: Scatter dry bait at the rate of $\frac{1}{4}$ lb. per 1,000 sq. ft. on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways, and window sills. On freshly limed floors, place treatment on feed sacks or on paper. Repeat procedures as necessary for practical fly control.

Caged Poultry and Turkeys on wire: Scatter dry bait at the rate of $\frac{1}{4}$ lb. per 100 sq. ft. on droppings beneath cages or porches. Repeat at weekly intervals.

Poultry Houses: Scatter dry bait at the rate of $\frac{1}{4}$ lb. per 100 sq. ft. where flies congregate but keep out of reach of birds. Apply to manure, on window sills, outside of penned enclosures and on floor of feed storage room. Repeat as necessary.

Dog Kennels: Scatter dry bait at the rate of $\frac{1}{4}$ lb. per 1,000 sq. ft. outside dog runways, window sills, ledges and other surfaces out of reach of animals. For caged units apply to dropping area. Repeat as necessary.

Garbage cans: Apply bait around and inside garbage can. Repeat as necessary.

For reducing populations of non resistant and resistant houseflies in and around outdoor living areas, picnic grounds, backyard areas, patios, loading docks, and outdoor latrines, parking areas, refuse areas around service stations, open air drive ins, ice cream stands, garbage collection and disposal areas: Use dry bait at $\frac{1}{2}$ lb. per 1,000 sq. ft. of area. Apply on ground or areas where flies congregate. For open air drive ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate.

CAUTION. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes, or clothing. Do not use inside the home. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed induce vomiting immediately and get medical attention. Atropine and 2-PAM are antidotal. Do not reuse container. Destroy when empty. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes the risk of use and/or handling of this material when it is used in a manner contrary to label instructions.

Koos, Inc. • Kenosha, Wisconsin 53140

