

HOLIDAY

20% Vapona[®]

(contains 2.0 lb. active ingredients per gallon)

ACTIVE INGREDIENTS	By Wt.
2-Ethoxypropyl isothiazyl phosphate	21.8%*
Related Compounds	1.6%*
Petroleum Hydrocarbons	65.6%
INERT INGREDIENTS	11.0%
	Total 100.0%

*Equivalent to 23.4% w/v VAPONA[®] insecticide

Net Weight:
32 oz. (1 qt.)



4500 Thirteenth Court / Kenosha, Wisconsin 53140

PRECAUTIONS IN USING

VAPONA[®] is a highly toxic insecticide. It is a contact poison and should be used only by trained personnel. It is not to be used on food crops, ornamentals, or in the presence of humans or animals. It is highly flammable and should be stored in a cool, dry place away from heat and open flames. It is highly volatile and should be used in a well-ventilated area. It is highly irritating to the eyes, nose, and throat. It is highly toxic to fish and aquatic life. It is highly toxic to birds and mammals. It is highly toxic to bees and other beneficial insects. It is highly toxic to cats and dogs. It is highly toxic to horses and ponies. It is highly toxic to swine and poultry. It is highly toxic to fish and aquatic life. It is highly toxic to birds and mammals. It is highly toxic to bees and other beneficial insects. It is highly toxic to cats and dogs. It is highly toxic to horses and ponies. It is highly toxic to swine and poultry. It is highly toxic to fish and aquatic life.

DANGER

VAPONA[®] is a highly toxic insecticide. It is a contact poison and should be used only by trained personnel. It is not to be used on food crops, ornamentals, or in the presence of humans or animals. It is highly flammable and should be stored in a cool, dry place away from heat and open flames. It is highly volatile and should be used in a well-ventilated area. It is highly irritating to the eyes, nose, and throat. It is highly toxic to fish and aquatic life. It is highly toxic to birds and mammals. It is highly toxic to bees and other beneficial insects. It is highly toxic to cats and dogs. It is highly toxic to horses and ponies. It is highly toxic to swine and poultry. It is highly toxic to fish and aquatic life.

DECONTAMINATE AND DESTROY USED CONTAINERS

Wash containers thoroughly with water before reuse. Do not reuse containers for food or drink. Destroy containers after use.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE & 2-PAM are the emergency antidotes for VAPONA[®] poisoning. Obtain them immediately by calling your local poison control center or hospital. **IN ALL CASES OF SUSPECTED POISONING**

INTERNAL: If the material has been swallowed, induce vomiting immediately by introducing a finger into the throat or by giving warm salt water. Follow with a glass of water. **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, remove the patient from the source of the hazard. Remove all contaminated clothing and shoes. Wash the exposed area with copious amounts of water. If the material gets into the eyes, wash immediately with running water for at least 15 minutes.

IF WARNING SYMPTOMS APPEAR: A person who has been exposed to VAPONA[®] should be kept under medical supervision. **WARNING SYMPTOMS:** Dizziness, headache, nausea, vomiting, diarrhea, and difficulty breathing.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Dizziness, headache, nausea, vomiting, diarrhea, and difficulty breathing. **TREATMENT:** Supportive care and administration of atropine and 2-pam.

DIRECTIONS FOR USE

ANIMAL BUILDINGS: Fogging animal buildings with VAPONA[®] is an effective method of controlling insect infestations. Fogging should be done in the absence of animals. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place.

bag Dumps Dog Kennels Fogging animal buildings with VAPONA[®] is an effective method of controlling insect infestations. Fogging should be done in the absence of animals. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place.

FOGGING ANIMAL BUILDINGS: Fogging animal buildings with VAPONA[®] is an effective method of controlling insect infestations. Fogging should be done in the absence of animals. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place.

CATTLE FEED LOTS STOCKYARDS HOLDING PENS CORRALS: Fogging animal buildings with VAPONA[®] is an effective method of controlling insect infestations. Fogging should be done in the absence of animals. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place.

CATTLE BEEF AND DAIRY: Fogging animal buildings with VAPONA[®] is an effective method of controlling insect infestations. Fogging should be done in the absence of animals. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place. Fogging should be done in a well-ventilated area. Fogging should be done in a cool, dry place.

KOGS, INC. • Kenosha, Wisconsin 53140

Kenosha, WI 53140

20% Vapona

(contains 2.0 lb. active ingredients per gallon)

Net Weight
32 oz (1 qt)



4500 Thirteenth Court / Kenosha, Wisconsin 53140

PRECAUTIONS IN USING

DANGER

DECONTAMINATE AND DESTROY USED CONTAINERS

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE & 2 PAM

INTERNAL

EXTERNAL

IF WARNING SYMPTOMS APPEAR

NOTE TO PHYSICIAN

WARNING SYMPTOMS

TREATMENT

DIRECTIONS FOR USE

ANIMAL BUILDINGS

Dairy Barns, Stables, Livestock Buildings, Kennels, Etc.

Large Ditches, Dog Kennels

FOGGING ANIMAL BUILDINGS

CATTLE FEED LOTS, HAYRACKS, FOGGING, ETC.

CATTLE, DEER AND DART